

Electricity and Magnetism

- Review for 8.02x Quiz #1
 - Electric Charge and Coulomb's Force
 - Electric Field and Field Lines
 - Superposition principle
 - E.S. Induction
 - Electric Dipole
 - Electric Flux and Gauss' Law
 - Electric Potential Energy and Electric Potential

Feb 18 2005

web.mit.edu/8.02x/www

Quiz

- Material as covered in class (this review)
- Evening Review in 54-100, 7PM, today
- Question(s) on demos (or similar setup)
 - all demos are fair game
- No question on LVPS/HVPS
 - but exp questions in future quizzes
- All you need is a pen
- You can bring a letter-size formula sheet
 - submitted together with quiz
- Quiz in room 26-100 **TUESDAY 10AM**

Feb 18 2005


web.mit.edu/8.02x/www

Electric Charge and Electrostatic Force

- New Property of Matter: **Electric Charge**
 - comes in **two flavors**: '+' and '-'
- Connected to **Electrostatic Force**
 - attractive (for '+-') or repulsive ('-', '+')
- Charge is **conserved**
- Charge is **quantized**
- Neutral: Equal amount of + and -

Feb 18 2005

web.mit.edu/8.02x/www



Feb 18 2005

web.mit.edu/8.02x/www

Coulomb's Law

$$\vec{F}_{12} = k \frac{Q_1 Q_2}{r_{21}^2} \hat{r}_{21} = \frac{1}{4\pi\epsilon_0} \frac{Q_1 Q_2}{r_{21}^2} \hat{r}_{21}$$


- Inverse square law ($F \sim 1/r^2$)
- Gives magnitude and direction of Force
- Attractive or repulsive depending on sign of $Q_1 Q_2$

Feb 18 2005

web.mit.edu/8.02x/www

Coulomb's Law

$$\vec{F}_{12} = k \frac{Q_1 Q_2}{r_{21}^2} \hat{r}_{21} = \frac{1}{4\pi\epsilon_0} \frac{Q_1 Q_2}{r_{21}^2} \hat{r}_{21}$$




Feb 18 2005

web.mit.edu/8.02x/www

Coulomb's Law

$$\vec{F}_{12} = k \frac{Q_1 Q_2}{r_{21}^2} \hat{r}_{21} = \frac{1}{4\pi\epsilon_0} \frac{Q_1 Q_2}{r_{21}^2} \hat{r}_{21}$$




Feb 18 2005

web.mit.edu/8.02s/www

Coulomb's Law

$$\vec{F}_{12} = k \frac{Q_1 Q_2}{r_{21}^2} \hat{r}_{21} = \frac{1}{4\pi\epsilon_0} \frac{Q_1 Q_2}{r_{21}^2} \hat{r}_{21}$$




$$\vec{F}_{12} = -\vec{F}_{21}$$

Feb 18 2005

web.mit.edu/8.02s/www

Superposition principle



- Note:
 - Total force is given by vector sum
 - Watch out for the charge signs
 - Use symmetry when possible

Feb 18 2005

web.mit.edu/8.02s/www

Superposition principle

- If we have many, many charges
 - Approximate with continuous distribution
- Replace sum with integral!

$$\vec{F}_{0,total} = \int d\vec{F}_0 = \int k \cdot \frac{Q_0 dQ}{r^2} \hat{r}$$

Feb 18 2005

web.mit.edu/8.02s/www

Electric Field

- New concept – Electric Field \vec{E}
- Charge Q gives rise to a Vector Field

$$\vec{E}(\vec{x}) = \vec{F}(\vec{x})/q$$

- \vec{E} is defined by strength and direction of force on small test charge q

Feb 18 2005

web.mit.edu/8.02s/www

The Electric Field

- Electric Field also exists if test charge q is not present
- The charge Q gives rise to a property of space itself – the Electric Field
- For more than one charge -> Superposition principle

Feb 18 2005

web.mit.edu/8.02s/www

Electric Field

- For a single charge

$$\vec{E} = k \frac{Q}{r^2} \hat{r}$$

Feb 18 2005


web.mit.edu/8.02s/www

Electric Field

- For a single charge

$$\vec{E} = k \frac{Q}{r^2} \hat{r}$$

- Visualize using Field Lines



Feb 18 2005

web.mit.edu/8.02s/www

Field Lines

- Rules for field lines


- Direction: Tangential to \vec{E} at each point
- Density: Shows magnitude of \mathbf{E}
- Field Lines never cross
- From positive to negative charge
 - i.e. show direction of force on a positive charge
- Far away: Everything looks like point charge

Feb 18 2005

web.mit.edu/8.02s/www

Example of Field Lines

- Field Lines for two unlike charges:




Feb 18 2005

web.mit.edu/8.02s/www

Holiday: Fundamentals of Physics

Electric Dipole




$$\text{Torque } \vec{\tau} = \vec{p} \times \vec{E}$$
$$\vec{p} = Q \vec{l} \text{ Dipolemoment}$$

Feb 18 2005

web.mit.edu/8.02s/www

Electrostatic Induction


- Approach neutral object with charged object



Feb 18 2005

web.mit.edu/8.02s/www

Electrostatic Induction



- Approach neutral object with charged object
- Induce charge separation (dipole)
- Force between charged and globally neutral object
 - if field is **non-uniform**

Feb 18 2005

web.mit.edu/8.02s/www

Electric Flux

- Electric Flux: $\Phi_E = \vec{E} \cdot \vec{A}$
- No 'substance' flowing
- Flux tells us how much field 'passes' through surface A

Feb 18 2005

web.mit.edu/8.02s/www

Electric Flux

- For 'complicated' surfaces and/or non-constant \vec{E} :
 - Use integral

$$\Phi_E = \int_A \vec{E} \cdot d\vec{A}$$

- Often, 'closed' surfaces


$$\Phi_E = \oint_A \vec{E} \cdot d\vec{A}$$

Feb 18 2005

web.mit.edu/8.02s/www

Electric Flux

- Example of closed surface: Box (no charge inside)



- Flux in (left) = -Flux out (right): $\Phi_E = 0$

Feb 18 2005

web.mit.edu/8.02s/www

Gauss' Law

- How are flux and charge connected?
- Charge **Q_{encl}** as source of flux through closed surface

$$\oint_A \vec{E} \cdot d\vec{A} = \frac{Q_{encl}}{\epsilon_0}$$

Feb 18 2005

web.mit.edu/8.02s/www

Gauss' Law

$$\oint_A \vec{E} \cdot d\vec{A} = \frac{Q_{encl}}{\epsilon_0}$$

- True for ANY closed surface around Q_{encl}
- Relates charges (cause) and field (effect)
- Coulombs Law follows from Gauss' Law

Feb 18 2005

web.mit.edu/8.02s/www

Gauss' Law

- Most uses of Gauss' Law rely on simple symmetries
 - Spherical symmetry
 - Cylinder symmetry
 - (infinite) plane
- and remember, $\mathbf{E} = \mathbf{0}$ inside conductors

Feb 18 2005

web.mit.edu/8.02s/www

Gauss' Law

- Different uses for Gauss' Law
 - Field \mathbf{E} -> Q_{encl} (e.g. conductor)
 - Q_{encl} -> Field \mathbf{E} (e.g. charged sphere)
- Proper choice of surface – use symmetries

$$\vec{E} \perp d\vec{A} \Rightarrow \vec{E} \cdot d\vec{A} = 0$$

$$\vec{E} \parallel d\vec{A} \Rightarrow \vec{E} \cdot d\vec{A} = E dA$$

$$E(r) = \text{const.} \Rightarrow \oint_A \vec{E} \cdot d\vec{A} = E \oint_A dA = EA$$


$$\vec{E} = 0 \Rightarrow \oint_A \vec{E} \cdot d\vec{A} = 0 = \frac{Q_{encl}}{\epsilon_0}$$

Feb 18 2005

web.mit.edu/8.02s/www

Hollow conducting Sphere

$$\vec{E} = 0 \Rightarrow \oint_A \vec{E} \cdot d\vec{A} = 0 = \frac{Q_{encl}}{\epsilon_0}$$




Feb 18 2005

web.mit.edu/8.02s/www

Gauss' Law

- Charge Sphere radius r_0 , charge Q , $r > r_0$

$$\begin{aligned} \frac{Q_{encl}}{\epsilon_0} &= \oint_{\text{sphere}} \vec{E} \cdot d\vec{A} = \\ &= \oint_{\text{sphere}} E dA = \\ &= E \oint_{\text{sphere}} dA = \\ &= E(4\pi r^2) \Rightarrow \\ E &= \frac{1}{4\pi\epsilon_0} \frac{Q_{encl}}{r^2} \end{aligned}$$




$\leftarrow Q_{encl} = Q$

Feb 18 2005

web.mit.edu/8.02s/www

Work and Potential Energy



Work: $W_{ba} = \int_a^b \vec{F} \cdot d\vec{l} = \int_a^b F dl \cos(\alpha)$

Conservative Force: $W_{ba} = -\Delta U = U(a) - U(b)$

Potential Energy

Feb 18 2005

web.mit.edu/8.02s/www

Electric Potential Energy

- Electric Force is conservative
 - all radial forces are conservative (e.g. Gravity)
- We can define Electric Potential Energy

$$W_{ba} = \int_a^b q\vec{E} \cdot d\vec{l} = U(a) - U(b) = -\Delta U$$

\uparrow
 \vec{F}

Feb 18 2005

web.mit.edu/8.02s/www

Electric Potential

- Electric Potential Energy proportional to q
- Define $\mathbf{V} = \mathbf{U}/q$

$$\frac{W_{ba}}{q} = \frac{U(a)}{q} - \frac{U(b)}{q} = V(a) - V(b) = -\Delta V$$

- Electric Potential \mathbf{V} :
 - Unit is Volt [V] = [J/C]

Feb 18 2005

web.mit.edu/8.02x/www

Electric Potential


- Note: because $\mathbf{V} = \mathbf{U}/q \rightarrow \mathbf{U} = \mathbf{V} q$
 - for a given \mathbf{V} : \mathbf{U} can be positive or negative, depending on sign of q
- \mathbf{V} : Work per unit charge to bring q from a to b

$$\frac{W_{ba}}{q} = \frac{U(a)}{q} - \frac{U(b)}{q} = V(a) - V(b) = -\Delta V$$


Feb 18 2005

web.mit.edu/8.02x/www

Example: Large parallel plates



Example: Large parallel plates



Conductors

- Conductor: Charges can move around (unlike insulator)
- $E = 0$ inside
 - otherwise charges would move
- No charges inside
 - Gauss
- E perpendicular to surface (close to surface)
 - otherwise charges on surface would move

Feb 18 2005

web.mit.edu/8.02x/www