

A GAUSSIAN FIXED POINT RANDOM WALK

YANG P. LIU, ASHWIN SAH, AND MEHTAAB SAWHNEY

ABSTRACT. In this note, we design a discrete random walk on the real line which takes steps $0, \pm 1$ (and one with steps in $\{\pm 1, 2\}$) where at least 96% of the signs are ± 1 in expectation, and which has $\mathcal{N}(0, 1)$ as a stationary distribution. As an immediate corollary, we obtain an online version of Banaszczyk’s discrepancy result for partial colorings and $\pm 1, 2$ signings. Additionally, we recover linear time algorithms for logarithmic bounds for the Komlós conjecture in an oblivious online setting.

1. INTRODUCTION

In the (oblivious) online vector discrepancy problem an adversary fixes vectors $\{v_i\}_{i \in [t]}$ in advance and the objective is to assign signs $\epsilon_i \in \{-1, 1\}$ based only on vectors v_1, \dots, v_i to maintain that $\|\sum_{i \leq t'} \epsilon_i v_i\|_\infty$ is small at all times $t' \in [t]$. Vector balancing includes a number of different problems in discrepancy theory including Spencer’s [17] work on set discrepancy. Spencer’s “six standard deviations suffice” result states that given vectors $v_1, \dots, v_n \in \{0, 1\}^n$ there exists a ± 1 -signing such that $\|\sum_{i \leq n} \epsilon_i v_i\|_\infty \leq 6\sqrt{n}$. Conjecturally, however, the restriction to $\{0, 1\}^n$ vectors can be relaxed to a norm condition. In particular, the Komlós conjecture states that given v_1, \dots, v_t , each of at most unit length, there exists a sequence of signs $\epsilon_1, \dots, \epsilon_t$ such that $\|\sum_{i \leq t} \epsilon_i v_i\|_\infty = O(1)$. Despite substantial effort, the Komlós conjecture is still open and the best known bounds due to Banaszczyk [4] give the existence of a sequence of signs so that $\epsilon_1, \dots, \epsilon_t$ such that $\|\sum_{i \leq t} \epsilon_i v_i\|_\infty = O(\sqrt{\min(\log n, \log t)})$. However, these original proofs were by their nature non-algorithmic.

More recent research in theoretical computer science has focused on developing algorithmic versions of these results starting with the Bansal [5] and Lovett-Meka [15] polynomial-time algorithms for Spencer’s [17] “six standard deviations suffice”. Since then, there have been several other constructive discrepancy minimization algorithms [16, 14, 6, 8, 7, 13]. Notably for our purposes, Bansal, Dadush, Garg [6] and Bansal, Dadush, Garg, Lovett [7] have made the work of Banaszczyk [4] algorithmic. However in all cases these algorithms require all vectors to be known at the start and hence do not extend to the online setting.

In the online setting, significant work has been devoted to the case where v_i are drawn from a fixed (and known) distribution \mathbf{p} supported on $[-1, 1]^n$. In the setting where \mathbf{p} is uniform on $[-1, 1]^n$, Bansal and Spencer [11] showed one can maintain $\max_{t' \leq t} \|\sum_{i \leq t'} \epsilon_i v_i\|_\infty \leq O(\sqrt{n} \log t)$. In the more general setting where \mathbf{p} is a general distribution supported on $[-1, 1]^n$, Aru, Narayanan, Scott, and Venkatesan [3] achieved a bound of $O_n(\sqrt{\log t})$ (where the implicit dependence on n is super-exponential) and Bansal, Jiang, Meka, Singla, and Sinha [9] (building on work of Bansal, Jiang, Singla, and Sinha [10]) achieved an ℓ_∞ guarantee of $O(\sqrt{n} \log(nt)^4)$.

In this work we focus on the online setting where the only guarantee is $\|v_i\|_2 \leq 1$. The only previous work in this oblivious online setting is the following result of Alweiss, the first author, and the third author [1].

Theorem 1.1 ([1, Theorem 1.1]). *For any vectors $v_1, v_2, \dots, v_t \in \mathbb{R}^n$ with $\|v_i\|_2 \leq 1$ for all $i \in [t]$, there exists an online algorithm $\text{BALANCE}(v_1, \dots, v_t, \delta)$ which maintains $\|\sum_{i \leq t'} \epsilon_i v_i\|_\infty = O(\log(nt/\delta))$ for all $t' \in [t]$ with probability at least $1 - \delta$.*

The proof in [1] relies on a coupling procedure which compares the distribution of $\sum_{i \leq t} \epsilon_i v_i$ to a Gaussian at each stage via a stochastic domination argument and then deduces the necessary tail bounds. In this work, we recover Theorem 1.1 (in fact with a slightly improved dependence) as well as the following corollary.

Corollary 1.2. *For any vectors $v_1, v_2, \dots, v_t \in \mathbb{R}^n$ with $\|v_i\|_2 \leq 1$ for all $i \in [t]$, there exists an online algorithm which assigns $\epsilon_i \in \{\pm 1, 2\}$ and maintains $\|\sum_{i \leq t'} \epsilon_i v_i\|_\infty = O(\sqrt{\log(nt/\delta)})$ for all $t' \in [t]$ with probability at least $1 - \delta$.*

This result essentially recovers the best known bound on the Komlós conjecture due to Banaszczyk [4] in an online algorithmic fashion, with the slight defect of requiring a +2-signing option. Furthermore due to the online nature of the algorithm, the algorithm will run in essentially input-sparsity time which is substantially faster than the Gram-Schmidt walk [7] which gives an algorithmic proof of the result of [4] (without the defect of requiring a +2-signing option).

Our results are based on the observation that there exists Markov chains on \mathbb{R} with transition steps of $0, \pm 1$ or $\pm 1, 2$ such that $\mathcal{N}(0, 1)$ is a stationary distribution (as well as $\mathcal{N}(0, \sigma^2)$ for appropriate values of σ). Note that no such walk exists for ± 1 steps as $\sum_{n \in \mathbb{Z}} (-1)^n e^{-n^2/2} \neq 0$ and therefore any ± 1 walk fails the natural “parity constraint” that the total mass on even integers is mapped to the odd integers and vice versa under one step.

The remainder of the paper is organized as follows. In Section 2 we construct the required Markov chain on \mathbb{R} with transition steps of $0, \pm 1$ such that $\mathcal{N}(0, \sigma^2)$ is a stationary distribution. In Section 3 we extend this to a walk with transition steps of $\pm 1, 2$ as long as $\sigma \geq 1$. Finally, in Section 4 we deduce the various algorithmic consequences.

1.1. Notation. Throughout this paper let $\mathcal{N}(\mu, \sigma^2)$ denote the Gaussian random variable with mean μ and variance σ^2 . Furthermore, let $\text{nnz}(\{v_i\}_{i \in S})$ denote the total number of non-zero entries of the vectors $\{v_i\}_{i \in S}$.

2. $0, \pm 1$ WALK

Definition 2.1. Given $\sigma > 0$ and $f \in [-1/2, 1/2]$, consider the following random walk on $f + \mathbb{Z}$. For $n \geq 1$ the state $n + f$ moves to $n + 1 + f$ with probability $p_\sigma(n + f)$ and to $n - 1 + f$ otherwise, and the state $-n + f$ moves to $-n - 1 + f$ with probability $p_\sigma(n - f)$ and to $-n + 1 + f$ otherwise. Finally, the state f moves to $1 + f$ with probability $p_\sigma(f)$, to state $-1 + f$ with probability $p_\sigma(-f)$, and stays at f with probability $r_\sigma(f)$. Here

$$p_\sigma(x) = \sum_{j \geq 1} (-1)^{j-1} \exp\left(-\frac{j^2 + 2xj}{2\sigma^2}\right)$$

$$r_\sigma(f) = \sum_{j=-\infty}^{\infty} (-1)^j \exp\left(-\frac{j^2 + 2fj}{2\sigma^2}\right)$$

for all $x \in \mathbb{R}$.

These series clearly absolutely converge. We prove that these indeed correspond to consistent probabilities giving a walk, and additionally show that this walk preserves the discrete Gaussian distribution on $f + \mathbb{Z}$ (i.e., $\mathcal{N}(0, \sigma^2)|_{f+\mathbb{Z}}$).

Lemma 2.2. *For $\sigma > 0$ and $f \in [-1/2, 1/2]$, we have that $p_\sigma(n \pm f) \in (0, 1)$ for all $n \geq 0$, that $p_\sigma(f) + r_\sigma(f) + p_\sigma(-f) = 1$, that $r_\sigma(f) \in [0, 1]$, and that furthermore*

$$r_\sigma(f) \leq e^{-\sigma^2}$$

if $\sigma \geq 1/2$. Additionally, $\mathcal{N}(0, \sigma^2)|_{f+\mathbb{Z}}$ is stationary under a step of random walk defined in Definition 2.1 with parameters σ, f .

Proof. First, note that $\exp(-(j^2 + 2xj)/\sigma^2)$ is strictly decreasing on integers $j \geq 1$ as long as $x \geq -1/2$. Therefore $p_\sigma(x)$ is given by an alternating series with strictly decreasing terms, and we immediately deduce

$$0 < p_\sigma(x) \leq \exp\left(-\frac{j^2 + 2xj}{2\sigma^2}\right) < 1.$$

Since $n + f, n - f \geq -1/2$ for $n \geq 0$, we see that $p_\sigma(n \pm f) \in (0, 1)$, as desired. Second, note that

$$p_\sigma(-f) + r_\sigma(f) + p_\sigma(f) = 1$$

holds as trivially everything except the $j = 0$ term of the sum for $r_\sigma(f)$ cancels. Third, we have for $u = \exp(-1/(2\sigma^2))$ and $v = \sqrt{-1} \exp(-f/(2\sigma^2))$ that $|u| < 1$ and $v \neq 0$, hence the Jacobi triple product identity (see [2] for a short but slick proof) yields

$$\begin{aligned} r_\sigma(f) &= \sum_{j=-\infty}^{\infty} u^{j^2} v^{2j} = \prod_{j=1}^{\infty} (1 - u^{2j})(1 + u^{2j-1}v^2)(1 + u^{2j-1}v^{-2}) \\ &= \prod_{j=1}^{\infty} (1 - e^{-j/\sigma^2})(1 - e^{-(2j+2f-1)/(2\sigma^2)})(1 - e^{-(2j-2f-1)/(2\sigma^2)}). \end{aligned} \quad (2.1)$$

Since $f \in [-1/2, 1/2]$ we see each term is nonnegative and clearly less than 1, so $r_\sigma(f) \in [0, 1]$ is immediate. Therefore we indeed have a well-defined walk. In fact, we see that

$$r_\sigma(f) \leq r_\sigma(0) \leq \prod_{j=1}^{\infty} (1 - e^{-j/\sigma^2})^3 \leq \prod_{j=1}^{\lfloor \sigma^2 \rfloor} (1 - e^{-j/\sigma^2})^3 \leq (1 - e^{-1})^{3\lfloor \sigma^2 \rfloor}.$$

This is at most $\exp(-\sigma^2)$ for $\sigma \geq 2$, and we can further numerically check that $r_\sigma(0) \leq \exp(-\sigma^2)$ for $\sigma \in [1/2, 2]$.

Now we show that this walk preserves $\mathcal{N}(0, \sigma^2)|_{f+\mathbb{Z}}$. Note that

$$1 - p_\sigma(x) = \sum_{j \geq 0} (-1)^j \exp\left(-\frac{j^2 + 2xj}{2\sigma^2}\right).$$

Therefore

$$\begin{aligned} &p_\sigma(x-1) \exp\left(-\frac{(x-1)^2}{2\sigma^2}\right) + (1 - p_\sigma(x+1)) \exp\left(-\frac{(x+1)^2}{2\sigma^2}\right) \\ &= \sum_{j \geq 1} (-1)^{j-1} \exp\left(-\frac{(j+x-1)^2}{2\sigma^2}\right) + \sum_{j \geq 0} (-1)^j \exp\left(-\frac{(j+x+1)^2}{2\sigma^2}\right) \\ &= \exp\left(-\frac{x^2}{2\sigma^2}\right). \end{aligned}$$

Since the pdf of $\mathcal{N}(0, \sigma^2)|_{f+\mathbb{Z}}$ at $n + f$ is proportional to $\exp(-(n + f)^2/(2\sigma^2))$, we find that the random walk preserves this distribution at $n + f$ for all $n \neq 0$ (applying the above equation at values $x = n \pm f$). Furthermore, the final distribution is clearly still supported on $f + \mathbb{Z}$, therefore the probability at $n = 0$ is also preserved as the total sum is 1. \square

We immediately derive a walk which preserves $\mathcal{N}(0, \sigma^2)$ by piecing together all $f \in [-1/2, 1/2]$. Let J_x^σ be the random variable defined by writing $x = n + f$, where $f \in [-1/2, 1/2]$, and then performing a step according to Definition 2.1.

Lemma 2.3. *If $Z = \mathcal{N}(0, \sigma^2)$ then $Z + J_Z^\sigma$ is distributed as $\mathcal{N}(0, \sigma^2)$.*

3. $\pm 1, 2$ WALK

We now consider a variant of the above random walk with discrete ± 1 and 2 steps. Recall the definition of $p_\sigma(x)$ and $r_\sigma(f)$ from earlier. We will require the following numerical estimate which is deferred to Appendix A.

Lemma 3.1. *If $\sigma \geq 1$ and $f \in [-1/2, 1/2]$ then*

$$p_\sigma(1+f) \geq r_\sigma(f) \exp\left(\frac{2f+1}{2\sigma^2}\right).$$

Remark. This inequality is immediate for large σ as the left uniformly tends to $1/2$ and the right uniformly decays to zero.

Definition 3.2. Given $\sigma \geq 1$ and $f \in [-1/2, 1/2]$, consider the following random walk on $f + \mathbb{Z}$. For $n \geq 2$ the state $n+f$ moves to $n+1+f$ with probability $p_\sigma(n+f)$ and to $n-1+f$ otherwise. For $n \geq 1$ the state $-n+f$ moves to $-n-1+f$ with probability $p_\sigma(n-f)$ and to $-n+1+f$ otherwise. The state f moves to $1+f$ with probability $p_\sigma(f)$, to state $-1+f$ with probability $p_\sigma(-f)$, and moves to $2+f$ with probability $r_\sigma(f)$. Finally, for $n = 1$ the state $1+f$ moves to $2+f$ with probability $p_\sigma(1+f) - r_\sigma(f) \exp((2f+1)/(2\sigma^2))$ and to f otherwise.

Lemma 3.3. *For $\sigma \geq 1$ and $f \in [-1/2, 1/2]$, we have that the walk in Definition 3.2 is well-defined, and that $\mathcal{N}(0, \sigma^2)|_{f+\mathbb{Z}}$ is stationary under a step of the walk with parameters σ, f .*

Proof. That all probabilities are valid follows from Lemma 2.2, except that we need to additionally verify

$$p_\sigma(1+f) \geq r_\sigma(f) \exp\left(\frac{2f+1}{2\sigma^2}\right).$$

This is precisely Lemma 3.1.

To verify that $\mathcal{N}(0, \sigma^2)|_{f+\mathbb{Z}}$ is preserved under the walk defined in Definition 3.2, recall that $\mathcal{N}(0, \sigma^2)|_{f+\mathbb{Z}}$ is preserved under walk defined in Definition 2.1 by Lemma 2.2. This walk only differs in its probabilities that f goes to $f, 2+f$ and that $1+f$ goes to $f, 2+f$. Therefore the probabilities at $n+f$ for $n \in \mathbb{Z} \setminus \{0, 2\}$ are correct. Since the probabilities sum to 1, it is enough to check the probability at $2+f$ is correct. It therefore suffices to show that

$$\begin{aligned} r_\sigma(f) \exp\left(-\frac{f^2}{2\sigma^2}\right) + \left(p_\sigma(1+f) - r_\sigma(f) \exp\left(\frac{2f+1}{2\sigma^2}\right)\right) \exp\left(-\frac{(1+f)^2}{2\sigma^2}\right) \\ + (1 - p_\sigma(3+f)) \exp\left(-\frac{(3+f)^2}{2\sigma^2}\right) = \exp\left(-\frac{(2+f)^2}{2\sigma^2}\right). \end{aligned}$$

We already verified in the proof of Lemma 2.2 that

$$p_\sigma(x-1) \exp\left(-\frac{(x-1)^2}{2\sigma^2}\right) + (1 - p_\sigma(x+1)) \exp\left(-\frac{(x+1)^2}{2\sigma^2}\right) = \exp\left(-\frac{x^2}{2\sigma^2}\right).$$

Plugging in $x = 2+f$ gives the desired identity, upon canceling the terms containing $r_\sigma(f)$. \square

Again, we immediately derive a walk which preserves $\mathcal{N}(0, \sigma^2)$ by piecing together all $f \in [-1/2, 1/2)$. Let R_x^σ be the random variable defined by writing $x = n+f$, where $f \in [-1/2, 1/2)$, and then performing a step according to Definition 2.1.

Lemma 3.4. *If $\sigma \geq 1$ and $Z = \mathcal{N}(0, \sigma^2)$ then $Z + R_Z^\sigma$ is distributed as $\mathcal{N}(0, \sigma^2)$.*

Algorithm 1: PARTIALCOLORING $_{\sigma}(v_1, \dots, v_t)$

```
1  $w_0 \leftarrow \mathcal{N}(0, \sigma^2 I_n)$ 
2 for  $1 \leq i \leq t$  do
3    $\sigma' \leftarrow \sigma / \|v_i\|_2$ 
4    $x' \leftarrow \langle w_{i-1}, v_i \rangle / \|v_i\|_2$ 
5    $w_i \leftarrow w_{i-1} + J_{x'}^{\sigma'} v_i.$ 
6  $w \leftarrow w_t - w_0$ 
```

Algorithm 2: BALANCING $_{\sigma}(v_1, \dots, v_t)$

```
1  $w_0 \leftarrow \mathcal{N}(0, \sigma^2 I_n)$ 
2 for  $1 \leq i \leq t$  do
3    $\sigma' \leftarrow \sigma / \|v_i\|_2$ 
4    $x' \leftarrow \langle w_{i-1}, v_i \rangle / \|v_i\|_2$ 
5    $w_i \leftarrow w_{i-1} + R_{x'}^{\sigma'} v_i.$ 
6  $w \leftarrow w_t - w_0$ 
```

4. ALGORITHMIC APPLICATIONS

We now derive a number of algorithmic consequences.

In both BALANCING $_{\sigma}$ and PARTIALCOLORING $_{\sigma}$, J and R are sampled independently every time. Additionally, note that BALANCING $_{\sigma}$ is only well-defined when $\sigma \geq 1$. Finally, we clearly see that PARTIALCOLORING $_{\sigma}$ assigns a sign of ± 1 to each given vector online, or chooses to omit it (a sign of 0), while BALANCING $_{\sigma}$ does the same except that the sign 2 is the additional alternative.

Our first algorithm application is a (weak version) of the partial coloring lemma.

Theorem 4.1. *Let $\|v_1\|_2, \dots, \|v_t\|_2 \leq 1$ and $\delta \in (0, 1/2)$. With probability at least $1 - \delta$ we have that $w_{\ell} - w_0$ in PARTIALCOLORING $_1(v_1, \dots, v_t)$ is $2\sqrt{2\log(2nt/\delta)}$ -bounded for all times $\ell \in [t]$. Furthermore, with probability at least $1 - \delta$ we have that $w_t - w_0$ is $2\sqrt{2\log(2n/\delta)}$ -bounded. Finally, at least 96.3% of vectors are used with probability $1 - \exp(-\Omega(t))$.*

Proof. By Lemma 2.3 we immediately see that $w_i \sim \mathcal{N}(0, \sigma^2 I_n)$ for all $i \in [t]$. The discrepancy results follow by trivial Gaussian estimates. For example, we see that the j th coordinate of w_{ℓ} is $\sqrt{2\log(2nt/\delta)}$ -bounded with probability at least $\delta/(2nt)$. Taking a union bound over $0 \leq \ell \leq t$ and $j \in [n]$ yields that w_0, \dots, w_t are bounded with probability at least $1 - \delta$. Therefore each difference is also bounded.

The fraction of vectors used being large follows from Chernoff's inequality and the fact that at every step, conditional on all previous choices, a vector is used with probability at least

$$\min_{f \in [-1/2, 1/2]} (1 - r_1(f)) \geq 0.9639. \quad \square$$

Our second algorithmic application recovers the online vector balancing results of Alweiss, the first author, and the third author [1, Theorems 1.1, 1.2].

Theorem 4.2. *Let $\|v_1\|_2, \dots, \|v_t\|_2 \leq 1$, $\delta \in (0, 1/2)$, and set $\sigma = \sqrt{\log(t/\delta)}$. With probability at least $1 - \delta$ we have that $w_{\ell} - w_0$ in PARTIALCOLORING $_{\sigma}(v_1, \dots, v_t)$ is $2\sqrt{2\log(t/\delta)\log(2nt/\delta)}$ -bounded for all times $\ell \in [t]$. Furthermore, with probability at least $1 - \delta$ we have that $w_t - w_0$ is $2\sqrt{2\log(t/\delta)\log(2n/\delta)}$ -bounded. Finally, all vectors are used with probability at least $1 - \delta$.*

Proof. The proof is essentially identical to that of Theorem 4.1. The only difference is that we see that at each step, a vector is not used with probability at most

$$\max_{f \in [-1/2, 1/2]} r_\sigma(f) \leq e^{-\sigma^2} = \frac{\delta}{t}$$

due to our choice of σ , by the inequality in Lemma 2.2. A union bound shows that all vectors are used with probability at least $1 - \delta$. \square

In fact, we can design an algorithm achieving the same bounds by using Algorithm 1 for any value of $\sigma \geq 1$ as follows. To do this, first run Algorithm 1, and then rerun Algorithm 1 on the vectors which were given a 0 sign until no vectors remain (note that this can still be done in an online manner). By Lemma 2.2, specifically $r_\sigma(f) \leq e^{-\sigma^2}$, this process will terminate with probability $1 - \delta$ in $O(\sigma^{-2} \log(t/\delta))$ rounds. Each run produces a random vector with variance $O(\sigma^2)$ in every coordinate, hence the total variance is $O(\log(t/\delta))$ per coordinate as desired.

Finally we recover an online version of Banaszczyk [4], except using $\pm 1, 2$ -signings. The proof is identical to that of Theorem 4.1 so we omit it.

Theorem 4.3. *Let $\|v_1\|_2, \dots, \|v_t\|_2 \leq 1$ and $\delta \in (0, 1/2)$. With probability at least $1 - \delta$ we have that $w_\ell - w_0$ in $\text{BALANCING}_1(v_1, \dots, v_t)$ is $2\sqrt{2 \log(2nt/\delta)}$ -bounded for all times $\ell \in [t]$. Furthermore, with probability at least $1 - \delta$ we have that $w_t - w_0$ is $2\sqrt{2 \log(2n/\delta)}$ -bounded.*

All three algorithmic procedures are online.

4.1. Computational details. In the previous section the above idealized algorithms ignored the cost of computing $r_\sigma(f)$ and $p_\sigma(n \pm f)$ to sufficient precision in order to be used for algorithmic purposes. The key claim is that one can approximate the above sums within δ in $\text{poly}(\log(\sigma/\delta))$ -time.

In order to do so first note that we can truncate the sums $p_\sigma(n \pm f)$ and $r_\sigma(f)$ to values of $j \geq 1$ where $(j^2 + 2(n \pm f)j)/(2\sigma^2) = O(\log(\sigma/\delta))$. We now note that

$$\left| e^x - \sum_{j=0}^m \frac{x^j}{j!} \right| \leq \frac{x^{m+1}}{(m+1)!} e^{\max(0, x)},$$

so taking $m = \Theta(\log(\sigma/\delta))$ gives a very good approximation to $\exp(-(j^2 + 2(n \pm f)j)/(2\sigma^2))$ in the range of terms considered. Now we can compute the desired sums by interpreting it as a sum of low degree (i.e. $O(\log(\sigma/\delta))$) polynomials on a sequence of integers, which can be evaluated quickly.

In the implementation of the algorithms above, at time t if we are given a vector shorter than $1/(2t^2)$, we deterministically add it but ignore it for the purposes of maintaining a Gaussian distribution. These vectors have total length at most 1, so contribute only $O(1)$ discrepancy in each coordinate. For the remaining vectors, we have $\sigma \leq 2t^2$. We thus can approximate the relevant probabilities to within $\delta/(2t^2)$ efficiently, and then sample appropriately. This will preserve the Gaussians in question up to total variation distance of at most

$$\sum_{t \geq 1} \frac{\delta}{2t^2} \leq \delta.$$

Therefore, the running time of all probability computations is $\text{poly}(\log(t/\delta))$ at time t . Thus the modified versions of the algorithms in Theorems 4.1 to 4.3 run in $O(t \text{poly}(\log(t/\delta)) + n + \text{nnz}(\{v_i\}_{i \in [t]}))$ time with discrepancy guarantees that are an absolute multiplicative factor worse. (The second term arises due to sampling the initial Gaussian point.) This running time essentially matches (up to logarithmic factors) the results of [1] and make progress towards input-sparsity time algorithms for discrepancy, a direction suggested by [12].

A variant of our algorithms which run in $O(t \text{poly}(\log(t/\delta)) + n \log t + \text{nnz}(\{v_i\}_{i \in [t]}))$ time is achieved by “disregarding vectors” at time t which are shorter than $1/(2t^2)$ (as above) and otherwise grouping vectors by length into dyadic scales and running the algorithms separately with independent randomness on each of the scales. Note that when vector lengths are forced to live in a dyadic scale then sampling an appropriate Gaussian leads us to compute the above probabilities only when $\sigma \in [1, 2]$ and hence direct evaluation of the series is efficient.

ACKNOWLEDGEMENTS

The authors thank Ryan Alweiss, Arun Jambulapati, Allen Liu, and Mark Sellke for earlier discussions on this topic. Furthermore we thank Ghaith Hiary for discussions regarding computing theta series.

REFERENCES

- [1] Ryan Alweiss, Yang P Liu, and Mehtaab Sawhney, *Discrepancy minimization via a self-balancing walk*, arXiv:2006.14009.
- [2] George E. Andrews, *A simple proof of Jacobi’s triple product identity*, Proc. Amer. Math. Soc. **16** (1965), 333–334.
- [3] Juhan Aru, Bhargav Narayanan, Alex Scott, and Ramarathnam Venkatesan, *Balancing sums of random vectors*, Discrete Anal. (2018), Paper No. 4, 17.
- [4] Wojciech Banaszczyk, *Balancing vectors and Gaussian measures of n -dimensional convex bodies*, Random Structures & Algorithms **12** (1998), 351–360.
- [5] Nikhil Bansal, *Constructive algorithms for discrepancy minimization*, 51th Annual IEEE Symposium on Foundations of Computer Science, FOCS 2010, October 23-26, 2010, Las Vegas, Nevada, USA, IEEE Computer Society, 2010, pp. 3–10.
- [6] Nikhil Bansal, Daniel Dadush, and Shashwat Garg, *An algorithm for Komlós conjecture matching Banaszczyk’s bound*, IEEE 57th Annual Symposium on Foundations of Computer Science, FOCS 2016, 9-11 October 2016, Hyatt Regency, New Brunswick, New Jersey, USA (Irit Dinur, ed.), IEEE Computer Society, 2016, pp. 788–799.
- [7] Nikhil Bansal, Daniel Dadush, Shashwat Garg, and Shachar Lovett, *The gram-schmidt walk: a cure for the Banaszczyk blues*, Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2018, Los Angeles, CA, USA, June 25-29, 2018 (Ilias Diakonikolas, David Kempe, and Monika Henzinger, eds.), ACM, 2018, pp. 587–597.
- [8] Nikhil Bansal and Shashwat Garg, *Algorithmic discrepancy beyond partial coloring*, Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing, 2017, pp. 914–926.
- [9] Nikhil Bansal, Haotian Jiang, Raghu Meka, Sahil Singla, and Makrand Sinha, *Online discrepancy minimization for stochastic arrivals*, Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms, SODA 2021, Virtual Conference, January 10 - 13, 2021 (Dániel Marx, ed.), SIAM, 2021, pp. 2842–2861.
- [10] Nikhil Bansal, Haotian Jiang, Sahil Singla, and Makrand Sinha, *Online vector balancing and geometric discrepancy*, Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing (New York, NY, USA), STOC 2020, Association for Computing Machinery, 2020, p. 1139–1152.
- [11] Nikhil Bansal and Joel H Spencer, *On-Line Balancing of Random Inputs*, arXiv:1903.06898.
- [12] Daniel Dadush, <https://homepages.cwi.nl/~dadush/workshop/discrepancy-ip/open-problems.html>.
- [13] Daniel Dadush, Aleksandar Nikolov, Kunal Talwar, and Nicole Tomczak-Jaegermann, *Balancing vectors in any norm*, 2018 IEEE 59th Annual Symposium on Foundations of Computer Science (FOCS), IEEE, 2018, pp. 1–10.
- [14] Ronen Eldan and Mohit Singh, *Efficient algorithms for discrepancy minimization in convex sets*, Random Struct. Algorithms **53** (2018), 289–307.
- [15] Shachar Lovett and Raghu Meka, *Constructive discrepancy minimization by walking on the edges*, SIAM J. Comput. **44** (2015), 1573–1582.
- [16] Thomas Rothvoss, *Constructive discrepancy minimization for convex sets*, SIAM Journal on Computing **46** (2017), 224–234.
- [17] Joel Spencer, *Six standard deviations suffice*, Trans. Amer. Math. Soc. **289** (1985), 679–706.

APPENDIX A. PROOF OF LEMMA 3.1

Proof of Lemma 3.1. First note $r_\sigma(f) \leq r_\sigma(0)$ and $r_\sigma(0) \leq r_1(0)$ follow immediately from the Jacobi triple product identity (2.1) and nonnegativity. Therefore it suffices to prove that

$$p_\sigma(1+f) \geq \exp(1/\sigma^2)r_1(0)$$

for all $\sigma \geq 1$ and $f \in [-1/2, 1/2]$.

First suppose that $\sigma \in [1, 2]$. Then since $p_\sigma(1+f)$ is an alternating series with decreasing terms,

$$p_\sigma(1+f) \geq \exp\left(-\frac{1+2(1+f)}{2\sigma^2}\right) - \exp\left(-\frac{2+2(1+f)}{\sigma^2}\right).$$

Fixing σ , the right has derivative

$$-\frac{1}{\sigma^2} \exp\left(-\frac{1+2(1+f)}{2\sigma^2}\right) + \frac{2}{\sigma^2} \exp\left(-\frac{2+2(1+f)}{\sigma^2}\right),$$

which we can check is positive for f underneath some cutoff and negative above this cutoff. Therefore the earlier expression is minimized over $f \in [-1/2, 1/2]$ at some $f \in \{\pm 1/2\}$. Then, numerical checking shows that for each case $f \in \{\pm 1/2\}$ the resulting expression is minimized on $\sigma \in \{1, 2\}$ for similar reasons. We find the true minimum is at $f = 1/2$ and $\sigma = 1$, which gives

$$p_\sigma(1+f) \geq 0.12 \geq er_1(0) \geq \exp(1/\sigma^2)r_1(0).$$

Now we suppose that $\sigma \geq 2$. Let $2k-1$ be the smallest odd integer larger than $\sigma-1-f$, which is clearly always a positive integer as $\sigma \geq 1$ and $f \leq 1/2$. We know that $t \mapsto \exp(-t^2/(2\sigma^2))$ is convex for $t \geq \sigma$, hence $t \mapsto \exp(-(t^2+2(1+f)t)/(2\sigma^2))$ is certainly convex and decreasing for $t \geq \sigma-1-f$. Therefore the difference between the values at j and $j+1$ is at least the difference between the values at $j+1$ and $j+2$ when $j \geq 2k-1$, yielding

$$\begin{aligned} p_\sigma(1+f) &= \sum_{j \geq 1} (-1)^{j-1} \exp\left(-\frac{j^2+2(1+f)j}{2\sigma^2}\right) \\ &\geq \sum_{j \geq 2k-1} (-1)^{j-1} \exp\left(-\frac{j^2+2(1+f)j}{2\sigma^2}\right) \\ &\geq \frac{1}{2} \left(\sum_{j \geq 2k-1} (-1)^{j-1} \exp\left(-\frac{j^2+2(1+f)j}{2\sigma^2}\right) + \sum_{j \geq 2k} (-1)^j \exp\left(-\frac{j^2+2(1+f)j}{2\sigma^2}\right) \right) \\ &= \frac{1}{2} \exp\left(-\frac{(2k-1)^2+2(1+f)(2k-1)}{2\sigma^2}\right) \\ &\geq \frac{1}{2} \exp\left(-\frac{(\sigma+1-f)^2+2(1+f)(\sigma+1-f)}{2\sigma^2}\right) \\ &\geq \frac{1}{2} \exp\left(-\frac{4\sigma^2+16\sigma+15}{8\sigma^2}\right) \geq \frac{1}{2} \exp(-71/32) \exp(1/\sigma^2) \\ &\geq 0.05 \exp(1/\sigma^2) \geq \exp(1/\sigma^2)r_1(0). \end{aligned} \quad \square$$

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA 94305, USA
Email address: yangpliu@stanford.edu

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA
Email address: {asah,msawhney}@mit.edu