

## Problems: Non-independent Variables

For question 3, I added  $z = 0$ . -HB

1. Find the total differential for  $w = zxe^y + xe^z + ye^z$ .

**Answer:**

$$\begin{aligned}dw &= ze^y dx + zxe^y dy + xe^y dz + e^z dx + xe^z dz + e^z dy + ye^z dz \\ &= (ze^y + e^z)dx + (zxe^y + e^z)dy + (xe^y + xe^z + ye^z)dz.\end{aligned}$$

2. With  $w$  as above, suppose we have  $x = t$ ,  $y = t^2$  and  $z = t^3$ . Write  $dw$  in terms of  $dt$ .

**Answer:** Here  $dx = dt$ ,  $dy = 2t dt$  and  $dz = 3t^2 dt$ . We do not substitute for  $x$ ,  $y$  and  $z$  because it does not greatly simplify the expression for  $dw$  and because in practice those values may be given or easily calculated from  $t$ .

$$dw = (ze^y + e^z)dt + (zxe^y + e^z)2t dt + (xe^y + xe^z + ye^z)3t^2 dt.$$

3. Now suppose  $w$  is as above,  $x^2y + y^2x = 1$ , and  $z = 0$ . Assuming  $x$  is the independent variable, find  $\frac{dw}{dx}$ .

**Answer:** The constraint  $x^2y + y^2x = 1$  becomes  $(2xy + y^2)dx + (x^2 + 2xy)dy = 0$ . Solving for  $dy$  in terms of  $x$ ,  $y$  and  $dx$  we get  $dy = \frac{2xy + y^2}{x^2 + 2xy} dx$ .

Using the equation for  $dw$  from (1) gives:

$$\begin{aligned}dw &= (ze^y + e^z)dx + (zxe^y + e^z)dy + (xe^y + xe^z + ye^z)dz \\ &= (0 + e^0)dx + (0 + e^0) \left( \frac{2xy + y^2}{x^2 + 2xy} dx \right) + 0 \\ &= dx + \frac{2xy + y^2}{x^2 + 2xy} dx \\ &= \frac{x^2 + 4xy + y^2}{x^2 + 2xy} dx.\end{aligned}$$

Thus,  $\frac{dw}{dx} = \frac{x^2 + 4xy + y^2}{x^2 + 2xy}$ .

MIT OpenCourseWare  
<http://ocw.mit.edu>

18.02SC Multivariable Calculus  
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.