

EVALUATING THE EFFECTIVENESS OF CHILD SAFETY SEATS AND SEAT BELTS IN PROTECTING CHILDREN FROM INJURY

JOSEPH J. DOYLE JR. and STEVEN D. LEVITT*

Young children are required to use child safety seats, and the age threshold at which children can legally graduate to seat belts has steadily increased. This article tests the relative effectiveness of child safety seats, lap-and-shoulder seat belts, and lap belts in preventing injuries among motor vehicle passengers aged 2–6 yr. We analyze three large, representative samples of crashes reported to police, as well as linked hospital data. We find no apparent difference in the two most serious injury categories for children in child safety seats versus lap-and-shoulder belts. Child safety seats provide a statistically significant 25% reduction in the least serious injury category. Lap belts are somewhat less effective than the two other types of restraints but far superior to riding unrestrained. (JEL 118)

I. INTRODUCTION

Motor vehicle crashes in the United States result in more than 45,000 deaths and an estimated 2.4 million disabling injuries each year (National Safety Council 2006). Restraint systems such as seat belts and airbags have been shown to dramatically limit the injuries sustained in a crash (Kahane 1986). For young children, all states currently require the use of child safety seats, and the minimum age and weight requirements to graduate to seat belts have been increasing over time (Insurance Institute for Highway Safety 2004). Since 2003, more than a dozen states raised these requirements.

Two important drawbacks of child safety seats are their cost and difficulty of correct installation. The 1997 Census of Manufactures reported that consumers in the United States spend more than \$300 million on roughly 4 million child safety seats each year. Meanwhile, it is estimated that 80% of child safety seats are incorrectly installed (U.S. Department of Transporta-

tion, 1996), reducing the safety benefits provided by child safety seats (Kahane 1986).

Seat belts offer a low-cost alternative to restrain children. All modern passenger vehicles come equipped with seat belts. Thus, there is no marginal cost to the consumer associated with their use. On the other hand, seat belts are primarily designed to fit adult passengers. Shoulder belts may fall improperly across the child's neck, and the lap belt may lie on the child's abdomen rather than across the hips, leading to possible abdominal injury and what is known as "seat belt syndrome."¹ In spite of these important drawbacks of seat belts for children, previous research has documented that children restrained by seat belts fare much better in crashes than unrestrained children (Hertz 1996; Johnston, Rivara, and Soderberg 1994; Partyka 1988).

There is a surprisingly limited body of research assessing the relative efficacy of child

1. See, for example, Kulowski and Rost (1956); Garrett and Braunstein (1962); Agran, Dunkle, and Winn (1987); Winston et al. (2000); Durbin, Arbogast, and Moll (2001); and Arbogast et al. (2002).

*We thank Paul Heaton and Jesse Shapiro for their comments. Ethan Lieber provided outstanding research assistance. Financial support was provided by the National Science Foundation and the Sherman Shapiro Research Fund.

Doyle: Associate Professor of Economics, Sloan School, Massachusetts Institute of Technology, Cambridge, MA 02142, Phone 617-452-3761, Fax 617-258-6855, E-mail jjdoyle@mit.edu

Levitt: Professor of Economics, Department of Economics, University of Chicago, 1126 East 59th Street, Chicago, IL 60637, Phone 773-834-1862, Fax 773-834-3040, E-mail slevitt@uchicago.edu

ABBREVIATIONS

CHSRA: Center for Health Systems
Research and Analysis
CODES: Crash Outcome Data Evaluation
System
GES: General Estimates Survey
NJDOT: New Jersey Department of
Transportation

safety seats and seat belts, and the existing studies come to very different conclusions. In a series of articles using a large sample of parental reports of injuries among children aged 4–7 yr, belt-positioning booster seats (the dominant form of child safety seat for this age group) have been found to reduce significant injuries by approximately 60% relative to seat belts (Durbin et al. 2003; Winston et al. 2000). Levitt (2005) uses police report data for crashes with at least one fatality and finds no statistically significant difference in fatalities or injuries between child safety seats and lap-and-shoulder belts for children in this age range.² That data set is far from ideal for studying injuries; however, as less than 2% of crashes with injuries involve a fatality.

In this article, we undertake the first comparison of the effectiveness of child safety seats and seat belts based on representative samples of all crashes reported to the police. We use three different data sets: (1) the General Estimates Survey (GES), a nationally representative sample of approximately 50,000 crashes each year for 16 yr; (2) New Jersey Department of Transportation (NJDOT) data covering all crashes in that state between 2001 and 2004; and (3) a Wisconsin data set that not only includes the universe of crashes with police reports in that state from 1994 to 2002 but also links these crashes to hospital discharge records. Using these data, we are able to exploit the wealth of information in police reports, as well as within-vehicle and within-accident variation in restraint use, to compare seat belts and child safety seats in preventing injury. The results suggest that lap-and-shoulder seat belts perform as well as child safety seats in preventing serious injury for children aged 2–6 yr. Safety seats are associated with a statistically significant reduction in the least serious injury category. According to our estimates, if every child wearing a lap-and-shoulder seat belt had instead been in a child safety seat, the number of injuries in this least serious category would be reduced by roughly 25%. Lap belts are somewhat less effective than the two other types of restraints but far superior to riding unrestrained.

The structure of the article is as follows. Section II describes the data sources and

provides some summary statistics. Section III describes the empirical strategy and main results. Section IV offers some conclusions and interprets the results in terms of the estimated benefits of injury reduction compared to the costs of increased safety seat use.

II. DATA AND SUMMARY STATISTICS

The data used are from three government collected, publicly available data sets: the GES, data from the NJDOT, and the Wisconsin Crash Outcome Data Evaluation System (CODES).

The GES is a nationally representative, stratified sample of all crashes reported to the police, collected annually by the National Highway Traffic Safety Administration for the period 1988–2004. Roughly 50,000 crashes are sampled each year. In all statistics and analyses reported in this article, the sample weights provided in the GES are used to make our estimates representative of the estimated 6.4 million crashes involving an injury or a significant property damage that occur annually.

NJDOT data include all motor vehicle crashes reported to the police in New Jersey over the period 2001–2004. One advantage of the New Jersey data is its large size, despite the fewer years available: more than 3.2 million people were involved in the 1.3 million police-reported crashes during these years.

Wisconsin CODES includes the universe of police accident reports from 1994 to 2002. These data have been linked to hospital discharge records by the University of Wisconsin's Center for Health Systems Research and Analysis (CHSRA). During these 9 yr, the system includes police reports for more than 3.2 million people and 2.1 million vehicles. CHSRA estimates that 80% of all crash-related hospitalizations were linked successfully, and the linkage rate does not vary systematically with the type of restraint used.³

For each of these data sets, the sample analyzed was restricted to children aged 2–6 yr. Few children younger than 2 yr use seat belts, and few children older than 6 yr use child safety seats in our data. We also restrict the

2. The study of Elliot et al. (2006), which combines the fatal accident data with another data set, the National Highway Traffic Safety Administration's National Automotive Sampling System Crashworthiness Data System, challenges the conclusions by Levitt (2005).

3. Emergency department admissions are available in Wisconsin CODES for 2002 and match the police-reported injury rates.

samples to children riding in passenger vehicles.⁴ Children riding in vehicle model years built prior to 1970 are excluded to ensure the presence of seat belts. Finally, approximately 10% of the data across the three sources are excluded due to missing values for restraint system used or the age of the passenger. The remaining samples include 46,205 observations in the GES (representing 5.5 million children applying the sampling weights); 74,971 observations in New Jersey; and 76,343 observations in Wisconsin.

Passengers are recorded in police reports as using no restraint, a child safety seat, a lap-and-shoulder belt, shoulder belt only, or lap belt only. No distinction is made in the data with respect to the precise type or model of child safety seat (e.g., backless vs. backed booster seats), and the data do not report whether a restraint was properly installed. We combine the small number of shoulder-only occupants with those wearing lap belts into a category we simply call “lap only,” though the results are similar regardless of how the shoulder-only occupants are categorized.

All three data sets use the KABC scale of injury severity, although different names are given to the A, B, and C injury categories in the data sets. In the GES and Wisconsin CODES, these four injury categories in order of decreasing severity are the following: fatal, incapacitating, nonincapacitating, and possible injury. In the NJDOT data, the four injury categories are labeled as: fatal, incapacitating, moderate, and complaint of pain. In all three data sets, the fatality rates are extremely low. Thus, in the analysis that follows, fatal and incapacitating injuries were combined into a single category of injury to provide a more precise estimate of the effectiveness of restraints in reducing the most severe injuries.

A. Summary Statistics

Summary statistics for the GES data set are reported in Table 1.⁵ Means for the full

4. The precise classification of vehicle types differs across the data sets. We include children riding in cars, minivans, and sport utility vehicles in the GES data; passenger cars and trucks in the Wisconsin data; and cars, vans, and sport utility vehicle/trucks in the New Jersey data.

5. For reasons of space, we do not report detailed summary statistics for the other two data sets. They are similar to the GES, with the exception that the New Jersey and Wisconsin data sets only include crashes from more recent years and thus are dominated by more recent vehicle model years, fewer children riding in the front seat, and fewer children unrestrained.

sample and each of the restraint types are included. Children in safety seats differ from other children in two important dimensions. Children riding in child safety seats are substantially younger on average than those using seat belts or riding unrestrained. Children in safety seats are also much less likely to be in the front seat than those using lap-and-shoulder belts, reflecting the availability of this type of restraint in front seats. Unrestrained children are also more likely to be found in the front seat. In all three data sets, we can reject equality of restraint use by seat position at the .01 level. Controlling for these factors is potentially important when estimating the effectiveness of restraints. Larger, older passengers have been shown to sustain greater injuries in crashes (Starnes 2005), and riding in the front seat is associated with a substantially elevated risk of injury (Kahane 2004; Starnes 2005).

Other than the differences in age and seat location, children in child safety seats and lap-and-shoulder belts have similar summary statistics, with some notable differences from those who are unrestrained or wearing lap belts. Lap belt-wearing children and those who are unrestrained are more likely to be riding in older model cars and thus will derive less benefit from recent vehicle safety advances (Kahane 2004). Children with no restraint also appear to be in vehicles with riskier drivers in general: the driver is less likely to be wearing a seat belt, more likely to be at fault in the crash, and more likely to be involved in a one-vehicle crash. Driver injuries are similar across the three types of restraints, but drivers of unrestrained children are found to suffer worse injuries, possibly because these drivers are more likely to be themselves unrestrained.

III. ESTIMATING THE RELATIVE EFFECTIVENESS OF CHILD SAFETY SEATS AND SEAT BELTS

Panel A of Table 2 presents the raw data on injury rates across restraint types for the three data sources. Rates of injury are substantially higher for unrestrained passengers. The differences between unrestrained passengers and those using any of the three restraints are highly statistically significant for all injury categories in all the data sets. Injury rates for children wearing seat belts are also somewhat higher compared to children in child safety seats.

TABLE 1
Summary Statistics: GES Sample

Characteristics	Full Sample		Means by Restraint Type			
	Mean	Standard Deviation	Safety Seat	Lap-and-Shoulder Belt	Lap-Only Belt	No Restraint
Child						
Age	3.91	1.39	2.92	4.42	4.28	4.12
Boy	0.51	0.50	0.51	0.50	0.50	0.51
Seat location						
Front	0.25	0.44	0.13	0.39	0.16	0.32
Back left	0.25	0.43	0.30	0.23	0.25	0.17
Back middle	0.14	0.35	0.17	0.06	0.23	0.17
Back right	0.27	0.45	0.34	0.25	0.28	0.19
Back other	0.005	0.07	0.002	0.001	0.006	0.018
Location unknown	0.08	0.26	0.06	0.07	0.08	0.13
Crash						
Front impact	0.42	0.49	0.42	0.43	0.41	0.44
Side impact	0.30	0.46	0.29	0.31	0.31	0.31
Rear impact	0.23	0.42	0.25	0.23	0.22	0.16
Other impact	0.003	0.06	0.003	0.004	0.003	0.003
Missing impact	0.04	0.21	0.03	0.03	0.05	0.09
One vehicle	0.12	0.32	0.12	0.10	0.12	0.18
Two vehicles	0.76	0.43	0.74	0.78	0.78	0.71
Three or more vehicles	0.12	0.33	0.13	0.12	0.11	0.11
Speed limit ≥ 55	0.17	0.38	0.18	0.16	0.17	0.21
Speed limit missing	0.13	0.34	0.16	0.10	0.15	0.11
Daytime	0.89	0.31	0.91	0.90	0.88	0.85
Evening	0.09	0.28	0.08	0.08	0.10	0.11
Early morning	0.01	0.12	0.01	0.01	0.02	0.03
Weekend	0.28	0.45	0.26	0.27	0.32	0.31
Vehicle						
Car	0.68	0.47	0.64	0.68	0.70	0.71
Minivan	0.13	0.34	0.16	0.13	0.11	0.10
Sport utility vehicle	0.19	0.39	0.20	0.19	0.18	0.19
Model year ≥ 1990	0.53	0.50	0.66	0.60	0.48	0.25
Driver						
Driver belted	0.91	0.29	0.48	0.96	0.98	0.96
Driver died	0.0009	0.031	0.0054	0.0004	0.0004	0.0002
Driver incapacitated	0.03	0.16	0.07	0.02	0.02	0.02
Driver nonincapacitated	0.06	0.23	0.10	0.05	0.05	0.05
Driver possibly injured	0.15	0.36	0.15	0.15	0.15	0.16
Driver no injury	0.76	0.43	0.65	0.78	0.78	0.77
Driver at fault	0.56	0.50	0.64	0.55	0.55	0.55
Observations	46,205		6,072	13,686	17,519	8,928

Note: The GES is a nationally representative sample of crashes from 1988 to 2004 available from National Highway Traffic Safety Administration, and sampling weights are applied to calculate nationally representative means.

The raw data on injury rates across restraint types may be misleading, however, given the differences across other observable characteristics. As noted earlier, one important factor to control for is seating position. Panel B of Table 2 also reports raw injury rates but

restricts the comparison to children riding in the back of the vehicle. The differences in outcomes between those in child seats and those in seat belts fall (and in some cases disappear completely) when the sample is restricted to back-seat passengers. The largest differences that

TABLE 2
Injuries Across Restraint Types

	Data: GES				Data: New Jersey				Data: Wisconsin			
	Child Seat	Lap-and-Shoulder Belt	Lap-Only Belt	No Restraint	Child Seat	Lap-and-Shoulder Belt	Lap-Only Belt	No Restraint	Child Seat	Lap-and-Shoulder Belt	Lap-Only Belt	No Restraint
Panel A: All children 2–6 yr old												
Fatal or incapacitating	.012	.015	.014	.070	.001	.001	.002	.013	.005	.006	.008	.039
Nonincapacitating	.034	.044	.048	.130	.019	.019	.023	.113	.026	.036	.038	.143
Possible injury	.077	.115	.107	.147	.096	.096	.116	.208	.060	.085	.086	.194
Observations	13,686	17,519	8,928	6,072	46,706	22,472	4,452	1,341	23,696	37,637	12,913	4,935
Panel B: Children aged 2–6 yr riding in backseat												
Fatal or incapacitating	.011	.011	.013	.061	.001	.001	.001	.017	.005	.005	.007	.033
Nonincapacitating	.032	.035	.042	.103	.012	.017	.023	.109	.025	0.028	.033	.111
Possible injury	.075	.108	.105	.142	.061	.092	.115	.201	.059	.079	.085	.180
Observations	11,976	10,857	7,509	4,192	45,592	19,481	4,177	1,105	20,686	23,685	11,220	3,385

Notes: The GES is a nationally representative sample of crashes from 1988 to 2004 available from National Highway Traffic Safety Administration, and sampling weights are applied to calculate nationally representative means. The NJDOT data contain all police-reported crashes in New Jersey from 2001 to 2004, and the injury severity categories are slightly differently labeled (fatal or incapacitating, moderate injury, and complaint of pain), although all three data sources use the KABC scale. The Wisconsin CODES data contain the universe of police-reported crashes in Wisconsin from 1994 to 2002.

persist are for the least serious injury category (“possible injury”). Riding unrestrained continues to be highly correlated with injury when the sample is limited to backseat passengers.

Seating position is, of course, just one of many factors that can contaminate measurement of the effectiveness of the various restraints. We use regression analysis to control for a wide range of potential confounding variables. In all cases, regressions are estimated using linear probability models. Probit models, evaluated at the sample mean, yield similar implied effects. The particular specification we estimate for child i involved in a crash in year t riding in a vehicle from model year m is as follows:

$$(1) \quad Y_i = \mathbf{R}_i\alpha + X_i\beta + \delta_t + \boldsymbol{\eta}_m + \varepsilon_i,$$

where Y is an indicator for a fatal or incapacitating injury, a nonincapacitating injury, or possible injury in three separate regressions. These injury categories are mutually exclusive and exhaustive. \mathbf{R} represents a vector of indicator variables describing the restraint type used, with child safety seats as the omitted category to test the difference in injury rates compared to a lap-and-shoulder belt, a lap belt, and a lack of restraint. X represents the control variables listed in Table 1: indicators for the age of the child, the principal point of impact in the crash, the time of day (early morning, daytime, and evening), whether the crash occurred on a weekend, the number of vehicles involved, whether the posted speed limit was 55 miles/h and above, and the type of vehicle (e.g., car, minivan, or sport utility vehicle/truck). Driver injuries, seat belt status, and the interaction between the driver injuries and seat belt status are also included as controls. δ_t is a vector of year dummies, and $\boldsymbol{\eta}_m$ is a vector of model year dummies. We also estimate models in which accident or vehicle fixed effects are included.

We present the results of this estimation on each of the three data sets in Tables 3–5. Table 3 reports the main results using the nationally representative GES database. Only the coefficients on the restraint types are included in the tables. Full regression results are available on request from the authors.⁶

6. The coefficients on these control variables are as expected. Riding in the front seat, for instance, greatly raises the risk of serious injury. The safest seat location, all else equal, is the back middle seat.

There are three sets of columns corresponding to the three outcomes: fatal or incapacitating injury, nonincapacitating injury, and possible injury. For each outcome, the results are presented for models with no controls, a second column that reports the results of models with seat location indicators, and a third column that includes the full set of controls.

The dependent variable in the first three columns of Table 3 is a fatal or incapacitating injury. As noted above, the omitted restraint category is child safety seats, so all estimates reported are relative to that category. Absent any controls (column 1), children wearing seat belts have somewhat higher rates of these injuries than those in child safety seats. Unrestrained children are much more likely to suffer fatal or incapacitating injuries. After controlling for seat position in column 2, the negative point estimate on lap-and-shoulder belts implies that children using this device actually experience a (small and statistically insignificant) reduction in fatal and incapacitating injuries relative to children in child safety seats. Controlling for seat position does not substantially affect the coefficients on lap-only belts or those who are unrestrained. Adding the full set of controls in column 3 leads the lap-and-shoulder belt coefficient to once again become positive (though statistically insignificant), doubles the coefficient on lap-only belts, and cuts in half the estimate for unrestrained passengers.⁷ Using our preferred estimates in column 3, relative to riding unrestrained, lap-and-shoulder belts provide 94% of the protection of child safety seats for fatal or incapacitating injuries.⁸ Lap-only belts

7. The driver injury categories exert the strongest influence among the controls. This reflects the fact that driver injuries are good proxies for the level of accident severity, though it may also reflect correlated measurement error. That is, measurement error in the child’s injury report and seat belt status may be correlated with measurement error in the driver injury report and driver seat belt status, especially since they are recorded by the same officer. Depending on whether and how this measurement error is related to the officer’s reporting of restraint types, it can bias the estimation in either direction. It is worth noting that omitting the driver injury measures does not systematically affect the estimates on restraint types.

8. Using a lap-and-shoulder belt increases these injuries by .0015 relative to child safety seats, compared to an increase of .0265 for riding unrestrained. The ratio of the coefficients on lap-and-shoulder belts and no restraint is approximately .06 (.0015/.0265), implying that lap-and-shoulder belts provide 94% of the benefits of child safety seats ($1 - .06 = .94$).

TABLE 3
Injury Comparison Across Restraint Types: GES

Dependent Variable	Fatal or Incapacitating			Nonincapacitating			Possible Injury		
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
Lap-and-shoulder belt	.0027 (.0011)*	-.0002 (.0012)	.0015 (.0013)	.0109 (.0020)**	.0026 (.0021)	.0014 (.0023)	.0378 (.0038)**	.0326 (.0039)**	.0183 (.0043)**
Lap-only belt	.0017 (.0014)	.0017 (.0014)	.0034 (.0014)*	.0143 (.0025)**	.0136 (.0025)**	.0093 (.0028)**	.0295 (.0046)**	.0287 (.0046)**	.0170 (.0049)**
No restraint	.0574 (.0033)**	.0560 (.0033)**	.0265 (.0030)**	.0967 (.0047)**	.0913 (.0047)**	.0558 (.0051)**	.0700 (.0060)**	.0646 (.0061)**	.0476 (.0071)**
Control for seat location	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Full controls	No	No	Yes	No	No	Yes	No	No	Yes
Observations	46,205	46,205	46,205	46,205	46,205	46,205	46,205	46,205	46,205
R ²	.0162	.0174	.2791	.017	.0207	.1239	.005	.0061	.1287
Mean of dependent variable	0.0201	0.0201	0.0201	0.0518	0.0518	0.0518	0.1058	0.1058	0.1058

Notes: Full controls include indicators for each age, seating position in the vehicle, principal point of impact, the number of vehicles in the crash, the type of vehicle, the time of the crash, whether it was on a weekend, if the speed limit was greater than 55 mph, year, and model year, as well as indicators for missing values for the seat location, speed limit, time of day, day of week, and model year. Driver controls include whether the driver was belted, the degree of injury sustained by the driver, interactions between the driver belt status and the driver injuries, and whether the driver was at fault. Excluded category is child safety seat. Sampling weights are applied to calculate nationally representative statistics.

*Significant at 5% level; **significant at 1% level.

TABLE 4
Injury Comparison Across Restraint Types: New Jersey

Dependent Variable	Fatal or Incapacitating			Moderate Injury			Report of Pain		
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
Lap-and-shoulder belt	.0004 (.0002)	.0001 (.0002)	.0002 (.0002)	.0065 (.0010)**	.0050 (.0010)**	.0043 (.0011)**	.0346 (.0023)**	.0317 (.0023)**	.0170 (.0023)**
Lap-only belt	.0018 (.0008)*	.0018 (.0007)*	.0017 (.0007)*	.0113 (.0023)**	.0112 (.0023)**	.0068 (.0023)**	.0546 (.0049)**	.0540 (.0050)**	.0291 (.0048)**
No restraint	.0128 (.0031)**	.0124 (.0032)**	.0108 (.0030)**	.1005 (.0086)	.0988 (.0086)**	.0768 (.0082)**	.1463 (.0111)**	.1428 (.0112)**	.0951 (.0106)**
Control for seat location	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Full controls	No	No	Yes	No	No	Yes	No	No	Yes
Observations	74,971	74,971	74,971	74,971	74,971	74,971	74,971	74,971	74,971
R ²	.0027	.0031	.0867	.0112	.0118	.1036	.0091	.0097	.1317
Mean of dependent variable	0.0011	0.0011	0.0011	0.0165	0.0165	0.0165	0.0780	0.0780	0.0780

Notes: Models with full controls include indicators for age, seating position in the vehicle, principal point of impact, the number of vehicles in the crash, the type of vehicle, the vehicle's model year, the time of the crash, whether it was on a weekend, if the speed limit was greater than 55 mph, year indicators, and model year indicators, as well as indicators for missing values for the seat location, model year, time of day, and missing speed limit. Driver controls include whether the driver was belted, the degree of injury sustained by the driver, and interactions between the driver belt status and the driver injuries. The omitted child restraint is child safety seat.

*Significant at the .05 level; **significant at the .01 level.

TABLE 5
Police-Reported Injury Comparison Across Restraint Types: Wisconsin

Dependent Variable	Fatal or Incapacitating			Nonincapacitating			Possible Injury		
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
Lap-and-shoulder belt	.0007 (.0006)	-.0004 (.0006)	-.0009 (.0007)	.0089 (.0014)**	.0031 (.0015)*	-.0027 (.0017)	.0245 (.0021)**	.0207 (.0022)**	.0097 (.0025)**
Lap-only belt	.0022 (.0009)*	.0025 (.0009)**	.0007 (.0010)	.0114 (.0020)**	.0120 (.0020)**	.0035 (.0022)	.0245 (.0030)**	.0250 (.0030)**	.0145 (.0032)**
No restraint	.0353 (.0030)**	.0347 (.0030)**	.0197 (.0028)**	.1197 (.0053)**	.1177 (.0054)**	.0795 (.0055)**	.1346 (.0061)**	.1341 (.0062)**	.0976 (.0066)**
Control for seat location	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Full controls	No	No	Yes	No	No	Yes	No	No	Yes
Observations	76,343	76,343	76,343	76,343	76,343	76,343	76,343	76,343	76,343
R ²	.0087	.0091	.1639	.0193	.0224	.1102	.0118	.0125	.1010
Mean of dependent variable	0.0079	0.0079	0.0079	0.0399	0.0399	0.0399	0.0843	0.0843	0.0843

Notes: Models with full controls include indicators for age, seating position in the vehicle, principal point of impact, the number of vehicles in the crash, the type of vehicle, the vehicle's model year, the time of the crash, whether it was on a weekend, if the speed limit was greater than 55 mph, year indicators, and model year indicators, as well as indicators for missing values for the speed limit, model year, and time of day. Driver controls include whether the driver was belted, the degree of injury sustained by the driver, interactions between the driver belt status and the driver injuries, and whether the driver was at fault. The omitted child restraint is child safety seat.

*Significant at the .05 level; **significant at the .01 level.

provide 87% of the protection of child safety seats in this injury category.

Columns 4–6 report parallel results for the second most severe injury category: nonincapacitating injuries. As more controls are added, the gap between child safety seats and the other categories falls. In the full specification, there is no statistically significant difference between children in lap-and-shoulder belts and those in child safety seats; for lap-only belts, the estimates are statistically significant. Relative to riding unrestrained, a lap-and-shoulder belt provides 97.5% of the benefit of a child safety seat, and a lap-only belt yields 83% of the benefit.

Columns 7–9 present results for the least serious injury category: possible injury. The patterns are similar to nonincapacitating injuries, except that child safety seats are now statistically significantly better than all the other devices. Both types of seat belts provide only about two-thirds of the benefits of child safety seats in preventing these least serious injuries.

Table 4 is identical in structure to Table 3, except that the results are based on the NJDOT data set rather than the GES. The patterns observed in the New Jersey data are quite similar to those in the GES. Adding controls tends to diminish the implied benefit of child safety seats. The point estimates suggest that lap-and-shoulder belts are 98% as effective as child safety seats for the most serious injuries (with the difference not statistically significant), 94% as effective for moderate injuries, and 83% as effective for the most minor injury category. In all three categories, lap-only belts are less effective than lap-and-shoulder belts.

Table 5 reports the same specifications but using the Wisconsin data. The results are similar to the other data sets, except that now with the full set of controls, lap-and-shoulder belts carry a point estimate that implies that they are (statistically insignificantly) better at preventing the two most serious injury types. The implied effectiveness of lap-and-shoulder belts relative to child safety seats in this sample is 105%, 103%, and 90% for the three injury categories. Lap-only belts are estimated to be 96%, 96%, and 85% as effective as child safety seats.

The consistency across the three data sets increases our confidence in the findings. There appears to be only minor differences between child safety seats and lap-and-shoulder belts in preventing fatal, incapacitating, and

nonincapacitating injuries. Only for the least serious category of injuries do we consistently observe substantial improvements from child safety seats relative to lap-and-shoulder belts, with these benefits ranging from 10 to 38% depending on the data set.

Because the Wisconsin crash data are linked to hospital discharge records, for that data set, we are able to explore a wider set of outcome variables, as shown in Table 6.

In the first two columns of the table, the dependent variable is an indicator equal to 1 if a child in a crash is admitted to the hospital and equal to 0 otherwise. Hospital admission represents inpatient care associated with more serious injuries, as opposed to just a visit to the emergency department. With or without controls, the point estimate on the lap-and-shoulder belt category is very close to 0 and statistically insignificant, implying no difference in hospitalization rates for children in child safety seats and those using lap-and-shoulder belts. The coefficient for lap belts is also small and statistically insignificant, implying that of every 700–800 children involved in a crash wearing a lap belt, only one fewer hospital admission would have occurred if those children were using child safety seats. In stark contrast, unrestrained passengers are five times more likely to be hospitalized compared to children using child safety seats or lap-and-shoulder belts.

The Wisconsin CODES data also provide measures of injury severity for those admitted to the hospital, summarized by the patient's length of stay and hospital charges.⁹ Greater treatment levels are positively correlated with police-reported injury severity and mortality.

The next four columns of Table 6 provide estimates of treatment differences across restraint types, conditional on hospital admission. The models are similar to Equation (1), with the natural logarithm of each treatment variable as the dependent variable.¹⁰ With

9. The Wisconsin data also allow us to look by location of the injury on the body. Lap-and-shoulder belts and child safety seats yield similar rates for injuries to the head, neck, and spine—those injuries most related to hospital charges and mortality. Lap belts are associated with slightly higher rates of head injury, and both types of belts were associated with slightly higher rates of leg injury.

10. The natural logarithm transformation provides more precise estimates, reflecting the skewed nature of hospital treatment data. Similar results are found for alternative functional forms. Results were also similar when the child's insurance status was included in these specifications.

TABLE 6
Hospital Outcomes Across Restraint Types: Wisconsin

Dependent Variable	Hospital Admission			Conditional on Hospital Admission		
	(1)	(2)	(3)	(4)	(5)	(6)
Lap-and-shoulder belt	-.0001 (.0005)	-.0002 (.0006)	-.2277 (.1390)	-.4407 (.1689)**	-.1914 (.1842)	-.3972 (.2161)
Lap-only belt	.0013 (.0007)	.0014 (.0008)	-.0249 (.1756)	-.2418 (.2005)	-.0350 (.2274)	-.1529 (.2525)
No restraint	.0249 (.0025)**	.0178 (.0024)**	-.0177 (.1415)	-.2234 (.2123)	-.0745 (.1770)	-.1580 (.2526)
Full controls	No	Yes	No	Yes	No	Yes
Observations	76,343	76,343	353	353	353	353
R ²	.0075	.0560	.0111	.2003	.0039	.2024
Mean of dependent variable	0.0046	0.0046	1.383	1.383	8.789	8.789

Notes: Models with full controls include indicators for age, seating position in the vehicle, principal point of impact, the number of vehicles in the crash, the type of vehicle, the vehicle's model year, the time of the crash, whether it was on a weekend, if the speed limit was greater than 55 mph, year indicators, and model year indicators, as well as indicators for missing values for the speed limit, model year, and time of day. Driver controls include whether the driver was belted, the degree of injury sustained by the driver, interactions between the driver belt status and the driver injuries, and whether the driver was at fault. The omitted child restraint is child safety seat.

*Significant at the .05 level; **significant at the .01 level.

the smaller sample size, these estimates become relatively imprecise, and the point estimates suggest that child safety seats are associated with worse injuries compared to seat belts. This (insignificant) difference is found for the comparison of safety seats and unrestrained passengers as well, which may reflect worse accidents required to induce a hospital admission for children in safety seats who are much less likely to be admitted into the hospital compared to children with no restraint.

A. Restraint Types and Accident Severity

One issue that arises when comparing injury rates is that the accident severity may vary across restraint types. Parents with a relatively high degree of risk aversion may be more likely to use safety seats *and* drive more safely. For example, while not a general result, de Meza and Webb (2001) and Jullien, Salanie, and Salanie (1999) suggest that individuals with a high degree of risk aversion may invest in self-protection as well as market insurance. Their results also suggest that careful driving behavior and restraint use may be complements. This would lead to an upward bias in the difference in injury rates between those wearing seat belts and those in safety seats. Indeed, as is noted earlier in Table 1, unrestrained children were more likely to be in vehicles where the driver is at fault and in a one-vehicle crash, though similar means were found for restrained passengers regardless of type. On the other hand, modes of self-protection may be substitutes, as in Peltzman (1975). Better protected riders may afford the driver the ability to take more risks on the road. This would lead to a downward bias in the above-mentioned results.

One approach to control for the accident severity is to include accident or vehicle fixed effects in the regression. These fixed effects will also absorb other unobserved sources of variation that may be contaminating the estimates, such as driving ability, road conditions, and the likelihood of the police officer to record particular types of injuries and restraint use. There are two drawbacks from using within-vehicle or within-accident variation in restraint use and injuries. First, less data can be used: only vehicles or accidents with more than one child involved provide variation in such a specification. Second, it relies on

children using different types of restraints in the same vehicle or accident. Within a vehicle, especially, there is relatively little variation in restraint use and the variation that is present is closely tied to the age of the children. Concerns about the endogeneity of restraint-type choice become heightened (e.g., if a parent puts one 4-yr-old in a child safety seat and another in a seat belt, that may reflect children of different weights, despite their similar ages). Concerns about coding errors in the data also increase: as almost all the variation is being removed from the data with vehicle fixed effects, a greater fraction of the variation that remains may be due to mistakes in the data.

With these caveats in mind, Table 7 reports the results of the fixed-effects specifications, where each panel is devoted to a particular data source. The three sets of columns again relate to the three different injury outcome variables. Specifications with accident and vehicle fixed effects are reported in the odd and even columns, respectively. All models include full controls, although most of those reported in Table 2 are absorbed by the fixed effects. Age, seat location, and sex are used in the vehicle fixed effects models, and indicators for model year, point of impact, vehicle type, and driver controls are also included in the accident fixed effects specification.

Because there are fewer observations involving multiple children in the same vehicle or accident, the results are relatively less precisely estimated. Nevertheless, the point estimates continue to show small differences for serious injuries. Estimated differences in the effectiveness of seat belts versus child safety seats for the least severe injury category are much smaller when accident or vehicle fixed effects are included for all three data sets. In the GES data, lap-and-shoulder belts have a coefficient of .0183 in the specification with full controls for the least severe injuries in Table 3, compared to a coefficient of .0108 when accident fixed effects are included—an increase in the relative effectiveness of lap-and-shoulder belts from 62% to 82%. For the New Jersey data, that same comparison yields .017 without accident fixed effects and .0067 with accident fixed effects (82% vs. 90%).

Table 8 explores the sensitivity of the basic results to various subgroups of the data. Because of space considerations, we limit the tabular presentation to the GES data set. The results are similar in the other data

sets. The three panels correspond to the injury categories; the columns correspond to the different subgroups.¹¹ The results are generally robust across the subgroups, though the smaller sample sizes result in less precise estimates. In terms of lap-and-shoulder belts, for none of the subgroups can we reject that the coefficients are equal to 0 for the two most serious injury categories. For possible injuries, we can reject equality for every subgroup, except the older age groups. For lap-only belts, about half of the coefficients across subgroups are statistically significant. Combining the information across injury severity levels, there does not appear to be strong evidence that the relative effectiveness of seat belts versus child safety seats differs sharply across the subgroups examined.¹²

IV. CONCLUSIONS

This study provides the first analysis of the relative effectiveness of seat belts and child safety seats in preventing injury based on representative samples of police-reported crash data. Our evidence suggests that lap-and-shoulder seat belts perform roughly as well as child safety seats in preventing serious injury for older children, although safety seats

11. Mean injury rates are given in the bottom row of each panel. Injury rates are found to increase with age, as found in previous research. Injuries are somewhat lower for vehicles with model years more recent than 1996, a period when more than 80% of children wearing a seat belt in the back left or back right of a vehicle used the lap-and-shoulder variety. Passenger cars have somewhat higher injury rates, likely attributable to their lower weight compared to minivans and sport utility vehicles.

12. Although not shown in tabular form, we have also investigated the relative effectiveness of the various restraints by the seriousness of the accident. To proxy for how serious the accident is, we used the sample of more than 300,000 passengers aged 7 and above in the GES to estimate specifications like those in Equation (1) with a dependent variable of fatal or incapacitating injury. Age fixed effects were used to control for age, and the coefficients on all nonage controls were used to predict (out of sample) the likelihood of serious injury for passengers aged 2–6 yr based on their characteristics, such as crash angle, seat location. We then divided individuals into five quintiles according to the predicted likelihood of serious injury. All the estimates were somewhat imprecise. The only statistically significant difference we observed between child safety seats and lap-and-shoulder belts for the two most serious injury categories was for the least severe quintile of crashes, where lap-and-shoulder belts are 80% as effective for both injury categories. The top four quartiles showed injury differences only for the least serious injury category.

TABLE 7
Accident or Vehicle Fixed Effects

Dependent Variable	Fatal or Incapacitating			Nonincapacitating			Possible Injury		
	(1)	(2)	(3)	(4)	(5)	(6)			
A. GES									
Lap-and-shoulder belt	.0011 (.0024)	.0022 (.0027)	.0081 (.0057)	.0036 (.0061)	.0108 (.0095)	.0098 (.0101)			
Lap-only belt	-.0007 (.0024)	-.0007 (.0027)	.0134 (.0065)*	.0111 (.0070)	.0139 (.0110)	.0153 (.0112)			
No restraint	.0148 (.0068)*	.0103 (.0070)	.0434 (.0126)**	.0369 (.0135)**	.0586 (.0180)**	.0536 (.0202)**			
Accident fixed effects	Yes	No	Yes	No	Yes	No			
Vehicle fixed effects	No	Yes	No	Yes	No	Yes			
Observations	17,779	16,111	17,779	16,111	17,779	16,111			
R ²	.8343	.8441	.6895	.7041	.7641	.7862			
Mean of dependent variable	0.0178	0.0183	0.0468	0.0476	0.0934	0.095			
B. New Jersey									
Lap-and-shoulder belt	.0005 (.0007)	.0009 (.0007)	.0009 (.0034)	.0011 (.0037)	.0067 (.0062)	.0090 (.0067)			
Lap-only belt	.0019 (.0013)	.0002 (.0004)	.0085 (.0068)	.0179 (.0071)*	-.0055 (.0100)	.0027 (.0109)			
No restraint	.0102 (.0067)	.0010 (.0008)	.0563 (.0243)*	.0916 (.0263)**	.0678 (.0264)*	.0622 (.0299)*			
Accident fixed effects	Yes	No	Yes	No	Yes	No			
Vehicle fixed effects	No	Yes	No	Yes	No	Yes			
Observations	27,236	23,695	27,236	23,695	27,236	23,695			
R ²	.6177	.6675	.6745	.7147	.7249	.7603			
Mean of dependent variable	0.0009	0.0008	0.0161	0.0161	0.0709	0.0696			
C. Wisconsin									
Lap-and-shoulder belt	-.0005 (.0015)	-.0012 (.0015)	-.0010 (.0042)	-.0036 (.0045)	.0014 (.0053)	.0016 (.0056)			
Lap-only belt	-.0010 (.0020)	-.0015 (.0018)	-.0056 (.0051)	-.0076 (.0055)	.0116 (.0068)	.0137 (.0072)			
No restraint	.0155 (.0055)**	.0156 (.0066)*	.0692 (.0149)**	.0768 (.0170)**	.0765 (.0181)**	.0697 (.0204)**			
Accident fixed effects	Yes	No	Yes	No	Yes	No			
Vehicle fixed effects	No	Yes	No	Yes	No	Yes			
Observations	30,974	28,310	30,974	28,310	30,974	28,310			
R ²	.7350	.7456	.6607	.6762	.7174	.7388			
Mean of dependent variable	0.0071	0.0356	0.0359	0.0072	0.0780	0.0784			

Notes: Data are restricted to accidents (vehicles) with more than one child present in the accident (vehicle) fixed effects specifications. All models are estimated with the full set of controls not absorbed by the fixed effects, including indicators for age, sex, and seat location in the vehicle fixed effects specifications, as well as model year, point of impact, vehicle type, and driver controls in the accident fixed effects specifications. The omitted child restraint is child safety seat. Sampling weights in the GES data are applied to calculate nationally representative statistics.

*Significant at the .05 level; **significant at the .01 level.

TABLE 8
Comparisons Across Subgroups in the GES

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
	Baseline	No Fault	Rear/Side Impact	Age 2	Age 4	Age 6	Backseat	Model Year ≥ 1997	Crash Year ≥ 1997	Car Only
A. Dependent variable: fatal or incapacitating										
Lap-and-shoulder belt	.0015 (.0013)	-.0009 (.0016)	.0014 (.0015)	.0043 (.0026)	-.0035 (.0025)	.0040 (.0036)	.0001 (.0013)	.0019 (.0019)	.0020 (.0015)	.0013 (.0016)
Lap-only belt	.0034 (.0014)*	.0039 (.0018)*	.0051 (.0018)**	.0009 (.0030)	-.0030 (.0029)	.0083 (.0042)*	.0032 (.0015)*	.0034 (.0030)	.0035 (.0020)	.0034 (.0018)
No restraint	.0265 (.0030)**	.0210 (.0038)**	.0187 (.0035)**	.0365 (.0070)**	.0178 (.0058)**	.0237 (.0071)**	.0244 (.0032)**	.0306 (.0108)**	.0310 (.0054)**	.0240 (.0035)**
Observations	46,205	18,975	23,251	9,880	9,779	8,094	34,534	8,698	24,572	31,199
R ²	.2791	.2674	.2893	.276	.3178	.339	.275	.3063	.2866	.2911
Mean of dependent variable	0.0201	0.0143	0.0153	0.0178	0.0214	0.0223	0.0169	0.0115	0.0178	0.0212
B. Dependent variable: nonincapacitating injury										
Lap-and-shoulder belt	.0014 (.0023)	-.0026 (.0033)	.0001 (.0030)	-.0019 (.0050)	.0038 (.0042)	-.0094 (.0117)	.0021 (.0025)	-.0010 (.0039)	.0024 (.0029)	.0000 (.0030)
Lap-only belt	.0093 (.0028)**	.0049 (.0037)	.0060 (.0037)	.0093 (.0070)	.0143 (.0051)**	.0080 (.0125)	.0046 (.0029)	.0097 (.0069)	.0118 (.0041)**	.0076 (.0035)*
No restraint	.0558 (.0051)**	.0363 (.0075)**	.0340 (.0065)**	.0487 (.0109)**	.0745 (.0112)**	.0381 (.0159)**	.0388 (.0054)**	.0504 (.0160)**	.0626 (.0081)**	.0550 (.0062)**
Observations	46,205	18,975	23,251	9,880	9,779	8,094	34,534	8,698	24,572	31,199
R ²	.1239	.1105	.1155	.1361	.1305	.1343	.1122	.1246	.1188	.1173
Mean of dependent variable	0.0518	0.0381	0.0399	0.0462	0.049	0.0598	0.043	0.032	0.0437	0.0552
C. Dependent variable: possible injury										
Lap-and-shoulder belt	.0183 (.0043)**	.0148 (.0070)*	.0150 (.0063)*	.0382 (.0092)**	.0002 (.0091)	.0234 (.0158)	.0186 (.0048)**	.0343 (.0088)**	.0205 (.0056)**	.0159 (.0055)**
Lap-only belt	.0170 (.0049)**	.0087 (.0081)	.0073 (.0072)	.0132 (.0099)	.0117 (.0104)	.0156 (.0175)	.0151 (.0053)**	.0034 (.0120)	.0131 (.0072)	.0151 (.0060)*
No restraint	.0476 (.0071)**	.0334 (.0122)**	.0317 (.0106)**	.0297 (.0132)*	.0464 (.0166)**	.0479 (.0228)*	.0419 (.0083)**	.0500 (.0204)*	.0414 (.0112)**	.0501 (.0088)**
Observations	46,205	18,975	23,251	9,880	9,779	8,094	34,534	8,698	24,572	31,199
R ²	.1287	.1219	.122	.1288	.135	.1348	.1263	.143	.1265	.1273
Mean of dependent variable	0.1058	0.1109	0.109	0.0833	0.1101	0.1236	0.0994	0.0911	0.1013	0.1137

Notes: Models use GES data, weighted to calculate nationally representative statistics. All models are estimated with the full set of controls. The omitted child restraint is child safety seat.

*Significant at the .05 level; **significant at the .01 level.

tend to be better at reducing less serious injuries. Lap belts had somewhat higher injury rates, while no restraint is associated with much larger injury rates. These results are robust to controls for accident severity, including within-vehicle and within-accident comparisons, as well as across age, seat location, and model year subgroups. Data on hospital admissions, length of stay, and hospital charges confirm the results based on police-reported injuries.

These comparisons across restraint types incorporate the way they are used (or misused) in practice. Because many child safety seats are, in actual use, improperly installed, our estimates are likely to understate the benefits associated with proper use of child safety seats. From a public policy perspective, however, understanding how safety devices work in practice, as opposed to under ideal circumstances, is of great importance.

These estimates of the benefits of child safety seats, which are based on samples reflecting all crashes reported to police in the areas represented, are far below those obtained in prior studies based on parent interviews. Even the upper bound estimates of the 95% confidence intervals that we estimate are much smaller than the benefits found in earlier studies using a survey-based approach.

Understanding why the results obtained from different sampling approaches vary so sharply is an important question. One possibility is that either parents (in the previous studies) or police (in the current study) systematically misreport restraint use or injuries. The congruence between police reports and hospital charges in the Wisconsin data provide external validation of the police codings. In addition, misreports to police would likely result in children being recorded as belted when they were actually unrestrained at the time of the crash. This type of misreporting would lead to an upward bias in the estimates presented here and thus is unlikely to explain the differing results. A second possibility is that we include a richer set of control variables than prior research. The three data sets we examine show large differences in injury rates in the raw data between children in child safety seats and those in seat belts. After controlling for a wide range of factors, however, the differences shrink dramatically or disappear.

The results allow an estimate of the cost-effectiveness of an investment in child safety

seats that are increasingly mandated for older children. In 1994, the Federal Highway Administration estimated monetary values associated with injuries on the KABC scale (Judycki 1994). The only category for which we observe systematic differences between child safety seats and lap-and-shoulder belts is the C category. Corrected for inflation between 1994 and 2004 using the Consumer Price Index, the current dollar value associated with this category of injury is \$24,130.¹³ This type of injury is roughly 25% higher among children using lap-and-shoulder belts relative to those in child safety seats. If every child aged 2–6 yr who was using a child safety seat instead switched to a lap-and-shoulder belt, the predicted increases in the number of C injuries in the United States in 2004 would have been approximately 3,000. The implied dollar value of these changes in injuries is \$72 million, below the roughly \$200 million spent annually by consumers on child safety seats for children aged 2–6 yr, but on the same order of magnitude. To the extent that there are extra benefits (albeit statistically insignificant ones in our analysis) of car seats in preventing more serious injuries, this estimate represents a lower bound on the safety value of car seats. This cost-benefit calculation is, of course, incomplete because it ignores other benefits (such as comfort for the child) and costs (parental time installing the seats) that accompany car seats. Nevertheless, the benefits of child safety seats in these three data sets are far less than implied by previous studies relying on parental survey data.

REFERENCES

- Agran, P., D. Dunkle, and D. Winn. "Injuries to a Sample of Seat Belted Children Evaluated and Treated in a Hospital Emergency Room." *Journal of Trauma*, 2, 1987, 58–64.
- Arbogast, K. B., D. R. Durbin, M. J. Kallan, R. A. Menon, A. E. Lincoln, and F. K. Winston. "The Role of Restraint and Seat Position in Pediatric Facial Fractures." *Journal of Trauma*, 52, 2002, 693–98.

13. The National Safety Council's Economics and Data Resource Center reports an estimated cost of these injuries due to wage and productivity losses, medical expenses, administrative expenses, motor vehicle damage, employers' uninsured costs, and a measure of the value of lost quality of life of \$22,900 in 2004 (National Safety Council 2004).

- de Meza, D., and D. C. Webb. "Advantageous Selection in Insurance Markets." *RAND Journal of Economics*, 32, 2001, 249–62.
- Durbin, D. R., K. B. Arbogast, and E. K. Moll. "Seat Belt Syndrome in Children: A Case Report and Review of the Literature." *Pediatric Emergency Care*, 17, 2001, 474–77.
- Durbin, D., M. Elliot, and F. Winston. "Belt-positioning Booster Seats and Reduction in Risk of Injury Among Children in Vehicle Crashes." *JAMA*, 289, 2003, 2835–40.
- Elliott, M. R., M. J. Kallan, D. R. Durbin, and F. K. Winston. "Effectiveness of Child Safety Seats vs Seat Belts in Reducing Risk for Death in Children in Passenger Vehicle Crashes." *Archives of Pediatrics & Adolescent Medicine*, 160, 2006, 617–21.
- Garrett, J. W., and P. W. Braunstein. "The Seat Belt Syndrome." *Journal of Trauma*, 2, 1962, 220–28.
- Hertz, E. *Revised Estimates of Child Restraint Effectiveness*. Washington, D.C.: National Highway Traffic Safety Administration, 1996.
- Insurance Institute for Highway Safety. "Child Restraint, Belt Laws as of July 2004." Accessed August 23, 2005. http://www.highwaysafety.org/safety_facts/state_laws/restrain.htm.
- Johnston, C., F. P. Rivara, and R. Soderberg. "Children in Car Crashes: Analysis of Data for Injury and Use of Restraints." *Pediatrics*, 93, 1994, 960–65.
- Judycki, D. C. *Motor Vehicle Accident Costs*. Washington, DC: Federal Highway Administration, 1994. Accessed August 23, 2005. <http://www.fhwa.dot.gov/legregs/directives/techadvst/75702.htm>.
- Jullien, B., B. Salanie, and S. Francois. "Should More Risk-Averse Agents Exert More Effort?" *Geneva Papers on Risk and Insurance Theory*, 1999, 24, 19–28.
- Kahane, C. *An Evaluation of Child Passenger Safety: The Effectiveness and Benefits of Safety Seats*. Washington, D.C.: National Highway Traffic Safety Administration, 1986. (Report no. DOT HS 806 890).
- . *Lives Saved by the Federal Motor Vehicle Safety Standards and Other Vehicle Safety Technologies*. Washington, D.C.: National Highway Traffic Safety Administration, 2004. (Report no. DOT HS 809 833).
- Kulowski, K., and W. Rost. "Intra-abdominal Injury from Safety Belts in Auto Accidents." *Archives of Surgery*, 73, 1956, 970–71.
- Levitt, S. "Evidence that Seat Belts are as Effective as Child Safety Seats in Preventing Death for Children Aged Two and Up." NBER Working Paper No. 11591, 2005.
- National Safety Council. "Estimating the Costs of Unintentional Injuries." 2004. Accessed August 23, 2005. <http://www.nsc.org/lrs/statinfo/estcost.htm>.
- National Safety Council. *Injury Facts*. Itasca, IL: National Safety Council, 2006.
- Partyka, S. *Lives Saved by Child Restraints from 1982 Through 1987*. Washington, D.C.: National Highway Traffic Safety Administration, 1988. (Report no. DOT HS 807 371).
- Peltzman, S. "The Effects of Automobile Safety Regulation." *Journal of Political Economy*, 83, 1975, 677–726.
- Starnes, M. *Child Passenger Fatalities and Injuries, Based on Restraint Use, Vehicle Type, Seat Position, and Number of Vehicles in the Crash*. Washington, D.C.: National Highway Traffic Safety Administration, 2005. (Report No. DOT HS 809 784).
- United States Department of Transportation. "Patterns of Misuse of Child Safety Seats." *Traffic Tech*, 133, 1996. Accessed July 15, 2008. <http://www.nhtsa.dot.gov/staticfiles/DOT/NHTSA/Communication%20&%20Consumer%20Information/Studies%20&%20Reports/1996/tt133.pdf>.
- Winston, F. K., D. R. Durbin, M. Kallan, and E. Moll. "The Danger of Premature Graduation to Seat Belts for Young Children." *Pediatrics*, 105, 2000, 1179–83.