

# On Krause's Multi-Agent Consensus Model With State-Dependent Connectivity

Vincent D. Blondel, Julien M. Hendrickx, and John N. Tsitsiklis, *Fellow, IEEE*

**Abstract**—We study a model of opinion dynamics introduced by Krause: each agent has an opinion represented by a real number, and updates its opinion by averaging all agent opinions that differ from its own by less than one. We give a new proof of convergence into clusters of agents, with all agents in the same cluster holding the same opinion. We then introduce a particular notion of equilibrium stability and provide lower bounds on the inter-cluster distances at a stable equilibrium. To better understand the behavior of the system when the number of agents is large, we also introduce and study a variant involving a continuum of agents, obtaining partial convergence results and lower bounds on inter-cluster distances, under some mild assumptions.

**Index Terms**—Consensus, decentralized control, multi-agent system, opinion dynamics.

## I. INTRODUCTION

THERE has been an increasing interest in recent years in the study of multi-agent systems where agents interact according to simple local rules, resulting in a possibly coordinated global behavior. In a prominent paradigm dating back to [11] and [29], each agent maintains a value which it updates by taking a linear, and usually convex combination of other agents' values; see e.g., [5], [17], [18], [26], [29], and [27], [28] for surveys. The interactions between agents are generally not all-to-all, but are described by an interconnection topology. In some applications, this topology is fixed, but several studies consider the more intriguing case of changing topologies. For example, in Vicsek's swarming model [31], animals are modeled as agents that move on the two-dimensional plane. All agents have the same speed but possibly different headings, and at each time-step they update their headings by averaging

the headings of those agents that are sufficiently close to them. When the topology depends on the combination of the agent states, as in Vicsek's model, an analysis that takes this dependence into account can be difficult. For this reason, the sequence of topologies is often treated as exogenous (see e.g., [4], [18], [26]), with a few notable exceptions [8], [9], [19]. For instance, the authors of [8] consider a variation of the model studied in [18], in which communications are all-to-all, but with the relative importance given by one agent to another weighted by the distance separating the agents. They provide conditions under which the agent headings converge to a common value and the distance between any two agents converges to a constant. The same authors relax the all-to-all assumption in [9], and study communications restricted to arbitrarily changing but connected topologies.

We consider here a simple discrete-time system involving endogenously changing topologies, and analyze it while taking explicitly into account the dependence of the topology on the system state. The discrete-agent model is as follows. There are  $n$  agents, and every agent  $i$  ( $i = 1, \dots, n$ ), maintains a real value  $x_i$ . These values are synchronously updated according to

$$x_i(t+1) = \frac{\sum_{j:|x_i(t)-x_j(t)|<1} x_j(t)}{\sum_{j:|x_i(t)-x_j(t)|<1} 1}. \quad (1)$$

Two agents  $i, j$  for which  $|x_i(t) - x_j(t)| < 1$  are said to be *neighbors* or *connected* (at time  $t$ ). Note that with this definition, an agent is always its own neighbor. Thus, in this model, each agent updates its value by computing the average of the values of its neighbors. In the sequel, we usually refer to the agent values as “opinions,” and sometimes as “positions.”

The model (1) was introduced by Krause [20] to capture the dynamics of opinion formation. Values represent opinions on some subject, and an agent considers another agent as “reasonable” if their opinions differ by less than 1<sup>1</sup>. Each agent thus updates its opinion by computing the average of the opinions it finds “reasonable”. This system is also sometimes referred to as the Hegselmann-Krause model, following [15]. It has been abundantly studied in the literature [20], [21], [23], [24], and displays some peculiar properties that have remained unexplained. For example, it has been experimentally observed that opinions initially uniformly distributed on an interval tend to converge to clusters of opinions separated by a distance slightly larger than 2, as shown in Fig. 1. In contrast, presently available results can only prove convergence to clusters separated by at least 1.

<sup>1</sup>In Krause's initial formulation, all opinions belong to  $[0, 1]$ , and an agent considers another one as reasonable if their opinions differ by less than a pre-defined parameter  $\epsilon$ .

Manuscript received July 11, 2008; revised July 15, 2008 and January 21, 2009. First published October 13, 2009; current version published November 04, 2009. This work was supported by the National Science Foundation under Grant ECCS-0701623, by the Concerted Research Action (ARC) “Large Graphs and Networks” of the French Community of Belgium and by the Belgian Programme on Interuniversity Attraction Poles initiated by the Belgian Federal Science Policy Office, and postdoctoral fellowships from the Belgian Fund for Scientific Research (F.R.S.-FNRS) and the Belgian American Education Foundation. (B.A.E.F). Recommended by Associate Editor M. Egerstedt.

V. D. Blondel is with the Department of Mathematical Engineering, Université catholique de Louvain, Louvain-la-Neuve B-1348, Belgium (e-mail: vincent.blondel@uclouvain.be).

J. M. Hendrickx is with the Laboratory for Information and Decision Systems, Massachusetts Institute of Technology, Cambridge, MA 02139 USA. He is also with the Department of Mathematical Engineering, Université catholique de Louvain, Louvain-la-Neuve B-1348, Belgium (e-mail: jm\_hend@mit.edu).

J. N. Tsitsiklis is with the Laboratory for Information and Decision Systems, Massachusetts Institute of Technology, Cambridge, MA 02139 USA (e-mail: jnt@mit.edu).

Color versions of one or more of the figures in this paper are available online at <http://ieeexplore.ieee.org>.

Digital Object Identifier 10.1109/TAC.2009.2031211




Fig. 1. Time evolution of 1000 agent opinions, according to the model (1). Initial opinions are either uniformly spaced (case (a)) or chosen at random (case (b)), on an interval of length 10. In both cases, opinions converge to limiting values (“clusters”) that are separated from each other by much more than the interaction radius, which was set to 1.

An explanation of the inter-cluster distances observed for this system, or a proof of a nontrivial lower bound is not available.

Inter-cluster distances larger than the interaction radius (which in our case was set to 1) have also been observed by Deffuant *et al.* [10] for a related stochastic model, often referred to as the Deffuant-Weisbuch model. In that model, two randomly selected agents update their opinions at any given time step. If their opinions differ by more than a certain threshold, their opinions remain unchanged; otherwise, each agent moves to a new opinion which is a weighted average of its previous opinion and that of the other agent. Thus, the Krause and Deffuant-Weisbuch models rely on the same idea of bounded confidence, but differ because one is stochastic while the other is deterministic. Besides, Krause’s model involves simultaneous interactions between potentially all agents, while the interactions in the Deffuant-Weisbuch model are pairwise. Despite these differences, the behavior of these two systems is similar, including inter-cluster distances significantly larger than the interaction radius. The behavior of the Deffuant-Weisbuch model—and in particular the final positions of the clusters—has also been studied by considering a continuous density approximating the discrete distribution of agents, and examining the partial differential equation describing the evolution of this density [2], [3]. Other models, involving either discrete or continuous time, and finitely or infinitely many agents, have also been proposed [1], [13], [30]. For a survey, see for example [25].

The model that we consider also has similarities with certain rendezvous algorithms (see, e.g., [22]) in which the objective is to have all agents meet at a single point. Agents are considered neighbors if their positions are within a given radius  $R$ . The update rules satisfy two conditions. First, when an agent moves, its new position is a convex combination of its previous position and the positions of its neighbors. Second, if two agents are neighbors, they remain neighbors after updating their positions. This ensures that an initially connected set of agents is never split into smaller groups, so that all agents can indeed converge to the same point.

In this paper, we start with a simple convergence proof for the model (1). We then introduce a particular notion of equilibrium stability, involving a robustness requirement when an equilibrium is perturbed by introducing an additional agent, and prove that an equilibrium is stable if and only if all inter-cluster distances are above a certain nontrivial lower bound. We observe

experimentally that the probability of converging to a stable equilibrium increases with the number of agents. To better understand the case of a large numbers of agents, we introduce and study a variation of the model, which involves a continuum of agents (the “continuous-agent” model). We give partial convergence results and provide a lower bound on the inter-cluster distances at equilibrium, under some regularity assumptions. We also show that for a large number of discrete agents, the behavior of the discrete-agent model indeed approximates the continuous-agent model.

Our continuous-agent model, first introduced in [6], is obtained by indexing the agents by a real number instead of an integer. It is equivalent to the so-called “discrete-time density based Hegselmann-Krause model” proposed independently in [25], which is in turn similar to a model presented in [13] in a continuous-time setup. Furthermore, our model can also be viewed as the limit, as the number of discrete opinions tends to infinity, of the “interactive Markov chain model” introduced by Lorenz [24]; in the latter model, there is a continuous distribution of agents, but the opinions take values in a discrete set.

We provide an analysis of the discrete-agent model (1) in Section II. We then consider the continuous-agent model in Section III. We study the relation between these two models in Section IV, and we end with concluding remarks and open questions, in Section V.

## II. THE DISCRETE-AGENT MODEL

### A. Basic Properties and Convergence

We begin with a presentation of certain basic properties of the discrete-agent model (1), most of which have already been proved in [15], [21], [23].

*Proposition 1 (Lemma 2 in [21]):* Let  $(x(t))$  be a sequence of vectors in  $\mathfrak{R}^n$  evolving according to (1). The order of opinions is preserved: if  $x_i(0) \leq x_j(0)$ , then  $x_i(t) \leq x_j(t)$  for all  $t$ .

*Proof:* We use induction. Suppose that  $x_i(t) \leq x_j(t)$ . Let  $N_i(t)$  be the set of agents connected to  $i$  and not to  $j$ ,  $N_j(t)$  the set of agents connected to  $j$  and not to  $i$ , and  $N_{ij}(t)$  the set of agents connected to both  $i$  and  $j$ , at time  $t$ . We assume here that these sets are nonempty, but our argument can easily be adapted if some of them are empty. For any  $k_1 \in N_i(t)$ ,  $k_2 \in N_{ij}(t)$ , and  $k_3 \in N_j(t)$ , we have  $x_{k_1}(t) \leq x_{k_2}(t) \leq x_{k_3}(t)$ . Therefore,  $\bar{x}_{N_i} \leq \bar{x}_{N_{ij}} \leq \bar{x}_{N_j}$ , where  $\bar{x}_{N_i}$ ,  $\bar{x}_{N_{ij}}$ ,  $\bar{x}_{N_j}$ , respectively, is the average of  $x_k(t)$  for  $k$  in the corresponding set. It follows from (1) that

$$x_i(t+1) = \frac{|N_{ij}|\bar{x}_{N_{ij}} + |N_i|\bar{x}_{N_i}}{|N_{ij}| + |N_i|} \leq \bar{x}_{N_{ij}},$$

and

$$x_j(t+1) = \frac{|N_{ij}|\bar{x}_{N_{ij}} + |N_j|\bar{x}_{N_j}}{|N_{ij}| + |N_j|} \geq \bar{x}_{N_{ij}}$$

where we use  $|A|$  to denote the cardinality of a set  $A$ . ■

In light of this result, we will assume in the sequel, without loss of generality, that the initial opinions are sorted: if  $i < j$  then  $x_i(t) \leq x_j(t)$ . The next Proposition follows immediately from the definition of the model.

*Proposition 2:* Let  $(x(t))$  be a sequence of vectors in  $\mathfrak{R}^n$  evolving according to (1), and such that  $x(0)$  is sorted, i.e., if

$i < j$ , then  $x_i(0) \leq x_j(0)$ . The smallest opinion  $x_1$  is nondecreasing with time, and the largest opinion  $x_n$  is nonincreasing with time. Moreover, if at some time the distance between two consecutive agent opinions  $x_i(t)$  and  $x_{i+1}(t)$  is larger than or equal to 1 it remains so for all subsequent times  $t' \geq t$ , so that the system can then be decomposed into two independent subsystems containing the agents  $1, \dots, i$ , and  $i+1, \dots, n$ , respectively.

Note that unlike other related models as the Deffuant-Weisbusch model [10] or the continuous-time model in [16], the average of the opinions is not necessarily preserved, and the ‘‘variance’’ (sum of squared differences from the average) may occasionally increase. See [16] for examples with three and eight agents respectively. The convergence of (1) has already been established in the literature (see [12], [23]), and is also easily deduced from the convergence results for the case of exogenously determined connectivity sequences (see e.g., [5], [17], [23], [26]), an approach that extends to the case of higher-dimensional opinions. We present here a simple alternative proof, which exploits the particular dynamics we are dealing with.

*Theorem 1:* If  $x(t)$  evolves according to (1), then for every  $i$ ,  $x_i(t)$  converges to a limit  $x_i^*$  in finite time. Moreover, for any  $i, j$ , we have either  $x_i^* = x_j^*$  or  $|x_i^* - x_j^*| \geq 1$ .

*Proof:* Since  $x(0)$  is assumed to be sorted, the opinion  $x_1$  is nondecreasing and bounded above by  $x_n(0)$ . As a result, it converges to a value  $x_1^*$ . Let  $p$  be the highest index for which  $x_p$  converges to  $x_1^*$ .

We claim that if  $p < n$ , there is a time  $t$  such that  $x_{p+1}(t) - x_p(t) \geq 1$ . Suppose, to obtain a contradiction, that the claim does not hold, i.e., that  $x_{p+1}(t) - x_p(t)$  is always smaller than 1. Fix some  $\epsilon > 0$  and a time after which the distance of  $x_i$  from  $x_1^*$ , for  $i = 1, \dots, p$ , is less than  $\epsilon$ . Since  $x_{p+1}$  does not converge to  $x_1^*$ , there is a further time at which  $x_{p+1}$  is larger than  $x_1^* + \delta$  for some  $\delta > 0$ . For such a time  $t$ ,  $x_p(t+1)$  is at least

$$\frac{1}{p+1} \left( \sum_{i=1}^{p+1} x_i(t) \right) \geq \frac{1}{p+1} (p(x_1^* - \epsilon) + (x_1^* + \delta))$$

which is larger than  $x_1^* + \epsilon$  if  $\epsilon$  is chosen sufficiently small. This however contradicts the requirement that  $x_p$  remain within  $\epsilon$  from  $x_1^*$ . This contradiction shows that there exists a time  $t$  at which  $x_{p+1}(t) - x_p(t) \geq 1$ . Subsequent to that time, using also Proposition 2,  $x_p$  cannot increase and  $x_{p+1}$  cannot decrease, so that the inequality  $x_{p+1} - x_p \geq 1$  continues to hold forever. In particular, agents  $1, \dots, p$  will no more interact with the remaining agents. Thus, if  $p < n$ , there will be some finite time after which the agents  $p+1, \dots, n$  behave as an independent system, to which we can apply the same argument. Continuing recursively, this establishes the convergence of all opinions to limiting values that are separated by at least 1.

It remains to prove that convergence takes place in finite time. Consider the set of agents converging to a particular limiting value. It follows from the argument above that there is a time after which none of them is connected to any agent outside that set. Moreover, since they converge to a common value, they eventually get sufficiently close so that they are all connected to each other. When this happens, they all compute the same average, reach the same opinion at the next time step, and keep

this opinion for all subsequent times. Thus, they converge in finite time. Finite time convergence for the entire systems follows because the number of agents is finite. ■

We will refer to the limiting values to which opinions converge as *clusters*. With some abuse of terminology, we will also refer to a set of agents whose opinions converge to a common value as a cluster.

It can be shown that the convergence time is bounded above by some constant  $c(n)$  that depends only on  $n$ . On the other hand, an upper bound that is independent of  $n$  is not possible, even if all agent opinions lie in the interval  $[0, L]$  for a fixed  $L$ . To see this, consider  $n$  agents, with  $n$  odd, one agent initially placed at 1, and  $(n-1)/2$  agents initially placed at 0.1 and 1.9. All agents will converge to a single cluster at 1, but the convergence time increases to infinity as  $n$  grows.

We note that the convergence result in Theorem 1 does not hold if we consider the same model but with a countable number of agents. Indeed, consider a countably infinite number of agents, all with positive initial opinions. Let  $m(y)$  be the number of agents having an initial opinion  $y$ . Suppose that  $\alpha \in (1/2, 1)$ , and consider an initial condition for which  $m(0) = 0$ ,  $m(\alpha) = 1$ ,  $m(\alpha(k+1)) = m(\alpha k) + 3m(\alpha(k-1))$  for every integer  $k > 1$ , and  $m(y) = 0$  for every other value of  $y$ . Then, the update rule (1) implies that  $x_i(t+1) = x_i(t) + \alpha/2$ , for every agent  $i$  and time  $t$ , and convergence fails to hold. A countable number of agents also admits equilibria where the limiting values are separated by less than 1. An example of such an equilibrium is obtained by considering one agent at every integer multiple of  $1/2$ .

We also note that equilibria in which clusters are separated by less than 1 become possible when opinions are elements of a manifold, instead of the real line. For example, suppose that opinions belong to  $[0, 2\pi)$  (identified with elements of the unit circle), and that two agents are neighbors if and only if  $|x_i - x_j \pmod{2\pi}| < 1$ . If every agent updates its angle by moving to the average of its neighbors' angles, it can be seen that an initial configuration with  $n$  agents located at angles  $2\pi k/n$ ,  $k = 0, \dots, n-1$ , is an equilibrium. Moreover, more complex equilibria also exist. Convergence has been experimentally observed for models of this type, but no proof is available.

## B. Experimental Observations

Theorem 1 states that opinions converge to clusters separated by at least 1. Since the smallest and largest opinions are nondecreasing and nonincreasing, respectively, it follows that opinions initially confined to an interval of length  $L$  can converge to at most  $\lceil L \rceil + 1$  clusters. It has however been observed in the literature that the distances between clusters are usually significantly larger than 1 (see [21], [24], and Fig. 1), resulting in a number of clusters that is significantly smaller than the upper bound of  $\lceil L \rceil + 1$ . To further study this phenomenon, we analyze below different experimental results, similar to those in [24].

Fig. 2 shows the dependence on  $L$  of the cluster number and positions, for the case of a large number of agents and initial opinions that are uniformly spaced on an interval of length  $L$ . Such incremental analyses also appear in the literature for various similar systems [2], [14], [24], [25]. We see that the cluster positions tend to change with  $L$  in a piecewise continuous (and sometimes linear) manner. The discontinuities correspond to the




Fig. 2. Locations of the different clusters at equilibrium, as a function of  $L$ , for  $5000L$  agents whose initial opinions are uniformly spaced on  $[0, L]$ , represented in terms of their distance from  $L/2$ . The dashed lines correspond to the endpoints 0 and  $L$  of the initial opinion distribution. Similar results are obtained if the initial opinions are chosen at random, with a uniform distribution.

emergence of new clusters, or to the splitting of a cluster into two smaller ones. The number of clusters tends to increase linearly with  $L$ , with a coefficient slightly smaller than  $1/2$ , corresponding to an inter-cluster distance slightly larger than 2. Note however that this evolution is more complex than it may appear: Irregularities in the distance between clusters and in their weights can be observed for growing  $L$ , as already noted in [24]. Besides, for larger scale simulations ( $L = 1000$ ,  $n = 10^6$ ), a small proportion of clusters take much larger or much smaller weights than the others, and some inter-cluster distances are as large as 4 or as small as 1.5. These irregularities could be inherent to the model, but may also be the result of the particular discretization chosen or of the accumulation of numerical errors in a discontinuous system.

Because no nontrivial lower bound is available to explain the observed inter-cluster distances in Krause's model, we start with three observations that can lead to some partial understanding. In fact, the last observation will lead us to a formal stability analysis, to be developed in the next subsection.

- a) We observe from Fig. 2 that the minimal value of  $L$  that leads to multiple clusters is approximately 5.1, while Theorem 1 only requires that this value be at least 1. This motivates us to address the question of whether a more accurate bound can be derived analytically. Suppose that there is an odd number of agents whose initial opinions are uniformly spaced on  $[0, L]$ . An explicit calculation shows that all opinions belong to an interval  $[1/2 - O(1/n), L - 1/2 + O(1/n)]$  after one iteration, and to an interval  $[11/12 - O(1/n), L - 11/12 + O(1/n)]$  after two iterations. Furthermore, by Proposition 2, all opinions must subsequently remain inside these intervals. On the other hand, note that with an odd number of agents, there is one agent that always stays at  $L/2$ . Thus, if all opinions eventually enter the interval  $(L/2 - 1, L/2 + 1)$ , then there can only be a single cluster. This implies that there will be a single cluster if  $L - 11/12 + O(1/n) < L/2 + 1$ , that is, if  $L < 23/6 - O(1/n) \simeq 3.833$ . This bound is smaller than the experimentally observed value of about 5.1. It can be

further improved by carrying out explicit calculations of the smallest position after a further number of iterations. Also, as long as the number of agents is sufficiently large, a similar analysis is possible if the number of agents is even, or in the presence of random initial opinions.

- b) When  $L$  is sufficiently large, Fig. 2 shows that the position of the leftmost clusters becomes independent of  $L$ . This can be explained by analyzing the propagation of information: at each iteration, an agent is only influenced by those opinions within distance 1 of its own, and its opinion is modified by less than 1. So, information is propagated by at most a distance 2 at every iteration. For the case of uniformly spaced initial opinions on  $[0, L]$ , with  $L$  large, the agents with initial opinions close to 0 behave, at least in the first iterations, as if opinions were initially distributed uniformly on  $[0, +\infty)$ . Moreover, once a group of opinions is separated from other opinions by more than 1, this group becomes decoupled. Therefore, if the agents with initial opinions close to 0 become separated from the remaining agents in finite time, their evolution under a uniform initial distribution on  $[0, L]$  for a sufficiently large  $L$  is the same as in the case of a uniform initial distribution on  $[0, +\infty)$ .

We performed simulations with initial opinions uniformly spaced on  $[0, +\infty)$ , as in [24]. We found that every agent eventually becomes connected with a finite number of agents and disconnected from the remaining agents. The groups formed then behave independently and converge to clusters. As shown in Fig. 3, the distances between two consecutive clusters are close to 2.2. These distances partially explain the evolution of the number of clusters (as a function of  $L$ ) shown in Fig. 2. However, a proof of these observed properties is not available, and it is unclear whether the successive inter-cluster distances possess some regularity or convergence properties.

- c) A last observation that leads to a better understanding of the size of the inter-cluster distances is the following. Suppose that  $L$  is just below the value at which two clusters are formed, and note the special nature of the resulting evolution, shown in Fig. 4. The system first converges to a "meta-stable state" in which there are two groups, separated by a distance slightly larger than 1, and which therefore do not interact directly with each other. The two groups are however slowly attracted by some isolated agents located in between; furthermore, these isolated agents are being pulled by both of these groups and remain at the weighted average of the opinions in the two groups. Eventually, the distance between the two groups becomes smaller than 1, the two groups start attracting each other directly, and merge into a single cluster. (This corresponds to one of the slow convergence phenomena observed in [24].) The initial convergence towards a two-cluster equilibrium is thus made impossible by the presence of a few agents in between. Moreover, the number of these isolated agents required to destabilize a meta-stable state can be arbitrarily small compared to the number of agents in the two groups. On the other hand, this phenomenon will not arise if the two clusters are separated by a sufficiently large distance. For example, if the distance




Fig. 3. Time evolution when the initial opinions are uniformly spaced on a semi-infinite interval, with a density of 100 per unit length. Groups of agents become separated from the remaining agents, and converge to clusters separated by approximately 2.2.




Fig. 4. Example of a temporary “meta-stable” state. Initially, two groups are formed that do not interact with each other, but they both interact with a small number of agents lying in between. As a result, the distance separating the two groups decreases slowly and eventually becomes smaller than 1. At that point, the groups attract each other directly and merge into a single cluster.

between the two groups is more than 2, no agent can be simultaneously connected to both groups. This suggests that, depending on the distance between clusters, some equilibria are stable with respect to the presence of a small number of additional agents, while some are not.

### C. Stability With Respect to a Perturbing Agent

In this section, we introduce a notion of equilibrium stability, motivated by the last observation in the preceding subsection. We first generalize the model (1), so that each agent  $i$  has an

associated weight  $w_i$  and updates its opinion according to the weighted discrete-agent model

$$x_i(t+1) = \frac{\sum_{j:|x_i(t)-x_j(t)|<1} w_j x_j(t)}{\sum_{j:|x_i(t)-x_j(t)|<1} w_j}. \quad (2)$$

It can be verified that the convergence results in Theorem 1 and the properties proved in Propositions 1 and 2 continue to hold. We will use the term *weight of a cluster* to refer to the sum of the weights of all agents in the cluster. Observe that if a number  $w$  of agents in system (1) have the same position, they behave as a single agent with weight  $w$  in the model (2). This correspondence can also be reversed, so that (2) can be viewed as a special case of (1), whenever the weights  $w_i$  are integer, or more generally, rational numbers.

Let  $\bar{x}$  be a vector of agent opinions at equilibrium. Suppose that we add a perturbing agent indexed by 0, with weight  $\delta$  and initial opinion  $\tilde{x}_0$ , that we let the system evolve again, until it converges to a new, perturbed equilibrium, and then remove the perturbing agent. The opinion vector  $\bar{x}'$  so obtained is again an equilibrium. We define  $\Delta_{\tilde{x}_0, \delta} = \sum_i w_i |\bar{x}_i - \bar{x}'_i|$ , which is a measure of the distance between the original and perturbed equilibria. We say that  $\bar{x}$  is *stable* if  $\sup_{\tilde{x}_0} \Delta_{\tilde{x}_0, \delta}$ , the supremum of distances between initial and perturbed equilibria caused by a perturbing agent of given weight  $\delta$ , converges to zero as  $\delta$  vanishes. Equivalently, an equilibrium is unstable if a substantial change in the equilibrium can be induced by a perturbing agent of arbitrarily small weight.

**Theorem 2:** An equilibrium is stable if and only if for any two clusters  $A$  and  $B$  with weights  $W_A$  and  $W_B$ , respectively, the following holds: either  $W_A = W_B$  and the inter-cluster distance is greater than or equal to 2; or  $W_A \neq W_B$  and the inter-cluster distance is *strictly* greater than  $1 + \min(W_A, W_B)/\max(W_A, W_B)$ . (Note that the two cases are consistent, except that the second involves a strict inequality.)

*Proof:* We start with an interpretation of the strict inequality in the statement of the theorem. Consider two clusters  $A$  and  $B$ , at positions  $x_A$  and  $x_B$ , and let  $m = (W_A x_A + W_B x_B)/(W_A + W_B)$ , which is their center of mass. Then, an easy calculation shows that

$$\begin{aligned} |x_A - x_B| > 1 + \frac{\min(W_A, W_B)}{\max(W_A, W_B)} \\ \text{if and only if} \\ \max\{|m - x_A|, |m - x_B|\} > 1. \end{aligned} \quad (3)$$

Suppose that an equilibrium  $\bar{x}_0$  satisfies the conditions in the theorem. We will show that  $\bar{x}_0$  is stable. Let us insert a perturbing agent of weight  $\delta$ . Note that since  $\bar{x}_0$  is an equilibrium, and therefore the clusters are at least 1 apart, the perturbing agent is connected to at most two clusters. If this agent is disconnected from all clusters, it has no influence, and  $\Delta_{\tilde{x}_0, \delta} = 0$ . If it is connected to exactly one cluster  $A$ , with position  $x_A$  and weight  $W_A$ , the system reaches a new equilibrium after one time step, where both the perturbing agent and the cluster have an opinion  $(\tilde{x}_0 \delta + x_A W_A)/(\delta + W_A)$ . Then

$$\Delta_{\tilde{x}_0, \delta} = |\tilde{x}_0 - x_A| \cdot \frac{\delta}{\delta + W_A} \leq \frac{\delta}{\delta + W_A}$$

which converges to 0 as  $\delta \rightarrow 0$ . Suppose finally that the perturbing agent is connected to two clusters  $A, B$ . This implies that the distance between these two clusters is less than 2, and since  $\tilde{x}_0$  satisfies the conditions in the theorem, it must be greater than  $1 + \min(W_A, W_B)/\max(W_A, W_B)$ . Therefore, using (3), the distance of one of these clusters from their center of mass  $m$  is greater than 1. The opinion of the perturbed agent after one iteration is within  $O(\delta)$  from  $m$ , while the two clusters only move by an  $O(\delta)$  amount. Since the original distance between one of the two clusters and  $m$  is greater than 1, it follows that after one iteration, and when  $\delta$  is sufficiently small, the distance of the perturbing agent from one of the clusters is greater than 1, which brings us back to the case considered earlier, and again implies that  $\Delta_{\tilde{x}_0, \delta}$  converges to zero as  $\delta$  decreases.

To prove the converse, we now suppose that the distance between two clusters  $A$  and  $B$ , at positions  $x_A$  and  $x_B$ , is less than 2, and also less than  $1 + \min(W_A, W_B)/\max(W_A, W_B)$ . Assuming without loss of generality that  $x_A < x_B$ , their center of mass  $m$  is in the interval  $(x_B - 1, x_A + 1)$ . Let us fix an  $\epsilon > 0$  such that  $(m - \epsilon, m + \epsilon) \subseteq (x_B - 1, x_A + 1)$ . Suppose that at some time  $t$  after the introduction of the perturbing agent we have

$$\tilde{x}_0(t) \in (m(t) - \epsilon, m(t) + \epsilon) \subseteq (x_B(t) - 1, x_A(t) + 1) \quad (4)$$

with  $x_B(t) - x_A(t) \geq 1$ , where  $\tilde{x}_0(t)$ ,  $x_A(t)$ ,  $x_B(t)$ , and  $m(t)$  represent the positions at time  $t$  of the perturbing agent, of the clusters  $A$  and  $B$ , and of their center of mass, respectively. One can easily verify that  $x_A(t+1) = x_A(t) + |\Theta(\delta)| > x_A(t)$ , and  $x_B(t+1) = x_B(t) - |\Theta(\delta)|$ , so that  $x_B(t+1) - x_A(t+1) < x_B(t) - x_A(t)$ , and  $(m(t+1) - \epsilon, m(t+1) + \epsilon) \subseteq (x_B(t+1) - 1, x_A(t+1) + 1)$ .

Moreover, observe that if  $\delta$  were 0, we would have  $\tilde{x}_0(t+1) = m(t)$ . For  $\delta \neq 0$ ,  $\tilde{x}_0(t+1)$  is close to  $m(t)$ , and we have  $\tilde{x}_0(t+1) = m(t) + O(\delta)$ . Since

$$\begin{aligned} m(t+1) &= \frac{W_A x_A(t+1) + W_B x_B(t+1)}{W_A + W_B} \\ &= m(t) + O(\delta) \end{aligned}$$

we obtain  $|m(t+1) - m(t)| = O(\delta)$ , and therefore  $\tilde{x}_0(t+1) \in (m(t+1) - \epsilon, m(t+1) + \epsilon)$ , as long as  $\delta$  is sufficiently small with respect to  $\epsilon$ .

We have shown that if  $\tilde{x}_0(0) = \tilde{x}_0$  is chosen so that the condition (4) is satisfied for  $t = 0$ , and if  $\delta$  is sufficiently small, the condition (4) remains satisfied as long as  $x_B(t) - x_A(t) \geq 1$ . The perturbing agent remains thus close to the center of mass, attracting both clusters, until at some time  $t^*$  we have  $x_B(t^*) - x_A(t^*) < 1$ . The two clusters then merge at the next time step. The result of this process is independent of the weight  $\delta$  of the perturbing agent, which proves that  $\bar{x}$  is not stable. Finally, a similar but slightly more complicated argument shows that  $\bar{x}$  is not stable when  $|x_A - x_B| = 1 + \min(W_A, W_B)/\max(W_A, W_B)$ , and  $|x_A - x_B| < 2$ . ■

Theorem 2 characterizes the stable equilibria in terms of a lower bound on the inter-cluster distances. It allows for inter-cluster distances at a stable equilibrium that are smaller than 2, provided that the clusters have different weights. This is consistent with experimental observations for certain initial opinion distributions, as shown in Fig. 5. On the other hand, for the




Fig. 5. Example of convergence to a stable equilibrium where the clusters are separated by less than 2. The initial distribution of opinions is obtained by taking 251 uniformly spaced opinions on  $[0, 2.5]$  and 500 uniformly opinions on  $[2.5, 3]$ . Opinions converge to two clusters with 153 and 598 agents, respectively, that are separated by a distance  $1.6138 > 1.2559 = 1 + 153/598$ . Similar results are obtained when larger number of agents are used, provided that the initial opinions are distributed in the same way, i.e. with a density on  $[2.5, 3]$  which is ten times larger than the density on  $[0, 2.5]$ .

frequently observed case of clusters with equal weights, stability requires the inter-cluster distances to be at least 2. Thus, this result comes close to a full explanation of the observed inter-cluster distances of about 2.2.

In general, there is no guarantee that the system (1) will converge to a stable equilibrium. (A trivial example is obtained by initializing the system at an unstable equilibrium, such as  $x_i(0) = -1/2$  for half of the agents and  $x_i(0) = 1/2$  for the other half). On the other hand, we have observed that for a given smooth distribution of initial opinions, and as the number of agents increases, we almost always obtain convergence to a stable equilibrium. This leads us to the following conjecture.

*Conjecture 1:* Suppose that the initial opinions are chosen randomly and independently according to a particular continuous and bounded probability density function (PDF) with connected support. Then, the probability of convergence to a stable equilibrium tends to 1, as the number of agents increases to infinity.

Besides the extensive numerical evidence (see e.g., Fig. 6), this conjecture is supported by the intuitive idea that if the number of agents is sufficiently large, whenever two groups of agents start forming two clusters, there will still be a small number of agents in between, whose presence will preclude convergence to an unstable equilibrium. The conjecture is also supported by Theorem 7 in Section III, which deals with a continuum of agents, together with the results in Section IV that provide a link between the discrete-agent and continuous-agent models.

### III. THE CONTINUOUS-AGENT MODEL

The discussion in the previous section indicates that much insight can be gained by focusing on the case of a large number of agents. This motivates us to consider a model involving a continuum of agents. We use the interval  $I = [0, 1]$  to index




Fig. 6. Time evolution of agent opinions, when initial opinions are drawn from a common PDF which is larger on the interval  $(2.5, 3)$  than on the interval  $(0, 2.5)$ . In (a), we have 501 agents and they converge to an unstable equilibrium: the clusters have respective weights 152 and 349, and their distance is  $1.399 < 1 + 152/349 \simeq 1.436$ . In (b), we have 5001 agents and they converge to a stable equilibrium: we see two clusters being formed originally, but they are eventually drawn together by a small number of agents in between.

the agents, and we consider opinions that are nonnegative and bounded above by a positive constant  $L$ . We denote by  $x_t(\alpha)$  the opinion of agent  $\alpha \in I$  at time  $t$ . We use  $X$  to denote the set of measurable functions  $x : I \rightarrow \mathfrak{R}$ , and  $X_L \subset X$  the set of measurable functions  $x : I \rightarrow [0, L]$ . The evolution of the opinions is described by

$$x_{t+1}(\alpha) = \frac{\int_{\beta: (\alpha, \beta) \in C_{x_t}} x_t(\beta) d\beta}{\int_{\beta: (\alpha, \beta) \in C_{x_t}} d\beta} \quad (5)$$

where  $C_x \subseteq I^2$  is defined for any  $x \in X$  by

$$C_x := \{(\alpha, \beta) \in I^2 : |x(\alpha) - x(\beta)| < 1\}.$$

If the denominator in (5) is zero, we use the convention  $x_{t+1}(\alpha) = x_t(\alpha)$ . However, since the set of agents  $\alpha$  for which this convention applies has zero measure, we can ignore such agents in the sequel. We assume that  $x_0 \in X_L$ . We then see that for every  $t > 0$ , we have  $x_t \in X_L$ , so that the dynamics are well-defined. In the sequel, we denote by  $\chi_x$  the indicator function of  $C_x$ , that is,  $\chi_x(\alpha, \beta) = 1$  if  $(\alpha, \beta) \in C_x$ , and  $\chi_x(\alpha, \beta) = 0$  otherwise.

We note that for the same reasons as in the discrete-agent model, if for some  $\alpha$  and  $\beta$  we have the relation  $x_t(\alpha) \leq x_t(\beta)$  or  $x_t(\alpha) = x_t(\beta)$  at some  $t$ , then the same relation continues to hold at all subsequent times. Furthermore, if  $x_0$  only takes a finite number of values, the continuous-agent model coincides with the weighted discrete-agent model (2), with the same range of initial opinions, and where each discrete agent's weight is set equal to the measure of the set of indices  $\alpha$  for which  $x_0(\alpha)$  takes the corresponding value.

In the remainder of this section, we will study the convergence properties of the continuous-agent model, and the inter-cluster distances at suitably defined stable equilibria.

### A. Operator Formalism

To analyze the continuous-agent model (5), it is convenient to introduce a few concepts, extending well known matrix and graph theoretic tools to the continuous case. By analogy with interaction graphs in discrete multi-agent systems, we define for

$x \in X$  the *adjacency operator*  $A_x$ , which maps the set  $X$  of measurable functions on  $I$  into itself, by letting

$$(A_x y)(\alpha) = \int \chi_x(\alpha, \beta) y(\beta) d\beta.$$

Applying this operator can be viewed as multiplying  $y$  by the “continuous adjacency matrix”  $\chi_x$ , and using an extension of the matrix product to the continuous case. We also define the *degree function*  $d_x : I \rightarrow \mathfrak{R}^+$ , representing the measure of the set of agents to which a particular agent is connected, by

$$d_x(\alpha) = \int \chi_x(\alpha, \beta) d\beta = (A_x \mathbf{1})(\alpha)$$

where  $\mathbf{1} : I \rightarrow \{1\}$  is the constant function that takes the value 1 for every  $\alpha \in I$ . Multiplying a function by the degree function can be viewed as applying an operator  $D_x : X \rightarrow X$  defined by

$$(D_x y)(\alpha) = d_x(\alpha) y(\alpha) = \int \chi_x(\alpha, \beta) y(\alpha) d\beta.$$

When  $d_x$  is positive everywhere, we can also define the operator  $D_x^{-1}$ , which multiplies a function by  $1/d_x$ . Finally, we define the *Laplacian operator*  $L_x = D_x - A_x$ . It follows directly from these definitions that  $L_x \mathbf{1} = 0$ , similar to what is known for the Laplacian matrix. In the sequel, we also use the scalar product  $\langle x, y \rangle = \int x(\alpha) y(\alpha) d\alpha$ . We now introduce two lemmas to ease the manipulation of these operators.

*Lemma 1:* The operators defined above are symmetric with respect to the scalar product: for any  $x, y, z \in X$ , we have  $\langle z, A_x y \rangle = \langle A_x z, y \rangle$ ,  $\langle z, D_x y \rangle = \langle D_x z, y \rangle$ , and  $\langle z, L_x y \rangle = \langle L_x z, y \rangle$ .

*Proof:* The result is trivial for  $D_x$ . For  $A_x$ , we have

$$\begin{aligned} \langle z, A_x y \rangle &= \int z(\alpha) \left( \int \chi_x(\alpha, \beta) y(\beta) d\beta \right) d\alpha \\ &= \int y(\beta) \left( \int \chi_x(\alpha, \beta) z(\alpha) d\alpha \right) d\beta. \end{aligned}$$

Since  $\chi_x(\alpha, \beta) = \chi_x(\beta, \alpha)$  for all  $\alpha, \beta$ , this implies  $\langle z, A_x y \rangle = \langle A_x z, y \rangle$ . By linearity, the result also holds for  $L_x$  and any other linear combination of those operators. ■

*Lemma 2:* For any  $x, y \in X$ , we have

$$\langle y, (D_x \pm A_x) y \rangle = \frac{1}{2} \int \chi_x(\alpha, \beta) (y(\alpha) \pm y(\beta))^2 d\alpha d\beta.$$

In particular,  $L_x = D_x - A_x$  is positive semi-definite.

*Proof:* From the definition of the operators, we have

$$\langle y, (D_x \pm A_x) y \rangle = \int \chi_x(\alpha, \beta) y(\alpha) (y(\alpha) \pm y(\beta)) d\alpha d\beta.$$

The right-hand side of this equality can be rewritten as

$$\begin{aligned} &\frac{1}{2} \left( \int \chi_x(\alpha, \beta) y(\alpha) (y(\alpha) \pm y(\beta)) d\alpha d\beta \right) \\ &+ \frac{1}{2} \left( \int \chi_x(\beta, \alpha) y(\beta) (y(\beta) \pm y(\alpha)) d\alpha d\beta \right). \end{aligned}$$

The symmetry of  $\chi_x$  then implies that  $\langle y, (D_x \pm A_x) y \rangle$  equals

$$\frac{1}{2} \int \chi_x(\alpha, \beta) (y(\alpha)^2 \pm 2y(\alpha)y(\beta) + y(\beta)^2) d\alpha d\beta$$

from which the results follows directly.  $\blacksquare$

The update (5) can be rewritten, more compactly, in the form

$$\begin{aligned} \Delta x_t &:= x_{t+1} - x_t = -D_x^{-1} L_{x_t} x_t, \text{ or} \\ D_{x_t} \Delta x_t &= -L_{x_t} x_t \end{aligned} \quad (6)$$

where the second notation is formally more general as it also holds on the possibly nonempty zero-measure set on which  $d_x = 0$ . We say that  $x_t \in X_L$  is a *fixed point* of the system if  $\Delta x_t = 0$  holds almost everywhere (a.e., for short), that is, except possibly on a zero-measure set. It follows from (6) that the set of fixed points is characterized by the equality  $L_x x = 0$ , a.e. One can easily see that the set of fixed points contains the set  $F := \{x \in X_L : x(\alpha) \neq x(\beta) \Rightarrow |x(\alpha) - x(\beta)| \geq 1\}$  of opinion functions taking a discrete number of values that are at least one apart. Let  $\bar{F}$  be the set of functions  $x \in X_L$  for which there exists  $s \in F$  such that  $s = x$ , a.e. We prove later that  $\bar{F}$  is exactly the set of solutions to  $L_x x = 0$ , a.e., and thus the set of fixed points of (6).

### B. Convergence

In this section we present some partial convergence results. In particular, we show that the change  $\Delta x_t$  of the opinion function decays to 0, and that  $x_t$  tends to the set of fixed points. We begin by proving the decay of a quantity related to  $\Delta x_t$ .

*Theorem 3:* For any initial condition of the system (6), we have

$$\sum_{t=0}^{\infty} \int \chi_{x_t}(\alpha, \beta) (\Delta x_t(\alpha) + \Delta x_t(\beta))^2 d\alpha d\beta < \infty.$$

*Proof:* We consider the nonnegative potential function  $V : X \rightarrow \mathbb{R}^+$  defined by

$$V(x) = \frac{1}{2} \int \min(1, (x(\alpha) - x(\beta))^2) d\alpha d\beta \geq 0 \quad (7)$$

and show that

$$V(x_{t+1}) - V(x_t) \leq -\langle \Delta x_t, (A_{x_t} + D_{x_t}) \Delta x_t \rangle$$

which by Lemma 2 implies the desired result.

We observe that for every  $x, y \in X$ , since  $\min(1, (y(\alpha) - y(\beta))^2)$  is smaller than or equal to both 1 and  $(y(\alpha) - y(\beta))^2$ , there holds

$$\begin{aligned} V(y) &\leq \frac{1}{2} \int_{C_x} (y(\alpha) - y(\beta))^2 d\alpha d\beta \\ &\quad + \frac{1}{2} \int_{I^2 \setminus C_x} 1 d\alpha d\beta \\ &= \langle y, L_x y \rangle + \frac{1}{2} |I^2 \setminus C_x| \end{aligned} \quad (8)$$

where Lemma 2 was used to obtain the last equability. For  $y = x$ , it follows from the definition of  $C_x$  that the above inequality is tight. In particular, the following two relations hold for any  $s$  and  $t$ :

$$\begin{aligned} V(x_t) &= \langle x_t, L_{x_t} x_t \rangle + \frac{1}{2} |I^2 \setminus C_{x_t}| \\ V(x_s) &\leq \langle x_s, L_{x_t} x_s \rangle + \frac{1}{2} |I^2 \setminus C_{x_t}|, \end{aligned}$$

Taking  $s = t + 1$ , we obtain

$$\begin{aligned} V(x_{t+1}) - V(x_t) &\leq \langle x_{t+1}, L_{x_t} x_{t+1} \rangle - \langle x_t, L_{x_t} x_t \rangle \\ &= 2\langle \Delta x_t, L_{x_t} x_t \rangle + \langle \Delta x_t, L_{x_t} \Delta x_t \rangle \end{aligned}$$

where we have used the symmetry of  $L_{x_t}$ . It follows from (6) that  $L_{x_t} x_t = -D_{x_t} \Delta x_t$ , so that

$$\begin{aligned} V(x_{t+1}) - V(x_t) &\leq -2\langle \Delta x_t, D_{x_t} x_t \rangle + \langle \Delta x_t, L_{x_t} x_t \rangle \\ &= -\langle \Delta x_t, (A_{x_t} + D_{x_t}) \Delta x_t \rangle \end{aligned}$$

since  $L_x = D_x - A_x$ .  $\blacksquare$

As will be seen below, this result implies the convergence of  $\Delta x_t$  to 0 in a suitable topology. We now show that  $L_x x$  is small only if  $x$  is close to  $F$ , the set of functions taking discrete values separated by at least 1. As a corollary, we then obtain the result that  $\bar{F}$  is exactly the set of fixed points, as also shown in [25]. The intuition behind the proof of these results parallels our proof of Theorem 1, and is as follows. Consider an agent  $\alpha$  with one of the smallest opinions  $x(\alpha)$ . If the change in  $x(\alpha)$  is small, its attraction by agents with larger opinions must be small, because almost no agents have an opinion smaller than  $x(\alpha)$ . Therefore, there must be very few agents with an opinion significantly larger than  $x(\alpha)$  that interact with  $\alpha$ , while there might be many of them who have an opinion close to  $x(\alpha)$ . In other words, possibly many agents have approximately the same opinion  $x(\alpha)$ , and very few agents have an opinion in the interval  $[x(\alpha) + \epsilon, x(\alpha) + 1]$ , so that  $x$  is close to a function in  $F$  in that zone. Take now an agent  $\alpha'$  with an opinion larger than  $x(\alpha) + 1 + \epsilon$ , and such that very few agents have an opinion in  $(x(\alpha) + 1 + \epsilon, x(\alpha'))$ . This agent interacts with very few agents having an opinion smaller than its own. Thus, if the change in such an agent's opinion is small, this implies that its attraction by agents having larger opinions is also small, and we can repeat the previous reasoning.

In order to provide a precise statement of the result, we associate an opinion function  $x$  with a measure that describes the distribution of opinions, and use a measure-theoretic formalism. For a measurable function  $x : I \rightarrow [0, L]$  (i.e.,  $x \in X_L$ ), and a measurable set  $S \subseteq [0, L]$ , we let  $\mu_x(S)$  be the Lebesgue measure of the set  $\{\alpha : x(\alpha) \in S\}$ . By convention, we let  $\mu(S) = 0$  if  $S \subseteq \mathbb{R} \setminus [0, L]$ . To avoid confusion with  $\mu$ , we use  $|S|$  to denote the standard Lebesgue measure of a set  $S$ . We also introduce a suitable topology on the set of opinion functions. We write  $x \leq_\mu \epsilon$  if  $|\{\alpha : x(\alpha) > \epsilon\}| \leq \epsilon$ . Similarly,  $x <_\mu \epsilon$  if  $|\{\alpha : x(\alpha) \geq \epsilon\}| < \epsilon$ , and  $x =_\mu 0$  if  $|\{\alpha : x(\alpha) \neq 0\}| = 0$ . We define the "ball"  $B_\mu(x, \epsilon)$  as the set  $\{y \in X_L : |x - y| <_\mu \epsilon\}$ . This allows us to define a corresponding notion of limit. We say that  $x_t \rightarrow_\mu y$  if for all  $\epsilon > 0$ , there is a  $t'$  such that for all  $t > t'$  we have  $x_t \in B_\mu(y, \epsilon)$ . We write  $x_t \rightarrow_\mu S$  for a set  $S$  if for all  $\epsilon > 0$ , there is a  $t'$  such that for all  $t > t'$ , there is a  $y \in S$  for which  $x_t \in B_\mu(y, \epsilon)$ .

The result below states that the distance between  $x \in X_L$  and  $F$  (the subset of  $X_L$  consisting of functions taking discrete values separated by at least 1) decreases to 0 (in a certain uniform sense) when  $L_x x \rightarrow_\mu 0$ . The proof, omitted for space reasons, is available in the appendix of [7] or in [16].

*Theorem 4:* For any  $\epsilon > 0$ , there exists a  $\delta > 0$  such that if  $|L_x x| <_\mu \delta$ , then there exists some  $s \in F$  with  $|x - s| <_\mu \epsilon$ . In particular, if  $L_x x =_\mu 0$ , then  $x \in \bar{F}$ .

The next theorem compiles our convergence results.

*Theorem 5:* Let  $(x_t)$  be a sequence of functions in  $X_L$  evolving according to the model (5), and let  $F$  be the set of functions taking discrete values separated by at least 1. Then  $(x_{t+1} - x_t) \rightarrow_{\mu} 0$  and  $x_t \rightarrow_{\mu} F$ . (In particular, periodic trajectories, other than fixed points, are not possible.) Furthermore,  $x$  is a fixed point of (5) if and only if  $x \in \bar{F}$ .

*Proof:* We begin by proving the convergence of  $\Delta x_t$ . Suppose that  $\Delta x_t = (x_{t+1} - x_t) \rightarrow_{\mu} 0$  does not hold. Then, there is an  $\epsilon > 0$  such that for arbitrarily large  $t$ , there is a set of measure at least  $\epsilon$  such that  $|\Delta x_t(\alpha)| > \epsilon$  for every  $\alpha$  in that set. Consider such a time  $t$ . Without loss of generality, assume that there is a set  $S \subseteq I$  of measure at least  $\epsilon/2$  on which  $\Delta x_t(\alpha) > \epsilon$ . (Otherwise, we can use a similar argument for the set on which  $\Delta x_t(\alpha) < -\epsilon$ .) Fix some  $L' > L$ . For  $i \in \{1, \dots, 2\lceil L' \rceil\}$ , let  $A_i \subset I$  be the set on which  $x_t \in [(i-1)/2, i/2]$ . For any  $i$  and for any  $\alpha, \beta \in A_i$ , there holds  $|x_t(\alpha) - x_t(\beta)| < 1$  and thus  $(\alpha, \beta) \in C_{x_t}$ . Therefore,  $A_i^2 \subseteq C_{x_t}$  for all  $i$ . Moreover, the sets  $A_i$  cover  $[0, 1]$ , so that  $\sum_{i=1}^{2\lceil L' \rceil} |A_i \cap S| \geq |S| \geq \epsilon/2$ . Thus, there exists some  $i^*$  such that  $|A_{i^*} \cap S| \geq \epsilon/(4\lceil L' \rceil)$ . We then have

$$\begin{aligned} & \int_{C_{x_t}} (\Delta x_t(\alpha) + \Delta x_t(\beta))^2 d\alpha d\beta \\ & \geq \int_{(A_{i^*} \cap S)^2} (\Delta x_t(\alpha) + \Delta x_t(\beta))^2 d\alpha d\beta \\ & \geq 4\epsilon^2 |A_{i^*} \cap S|^2 \geq \frac{\epsilon^4}{4\lceil L' \rceil^2}. \end{aligned}$$

Thus, if  $\Delta x_t \rightarrow_{\mu} 0$  does not hold, then  $\int_{(\alpha, \beta) \in C_{x_t}} (\Delta x_t(\alpha) + \Delta x_t(\beta))^2$  does not decay to 0, which contradicts Theorem 3. We conclude that  $\Delta x_t \rightarrow_{\mu} 0$ . Using also (6) and the fact  $d_{x_t}(\alpha) \leq 1$ , we obtain  $L_{x_t} x_t \rightarrow_{\mu} 0$ . Theorem 4 then implies that  $x_t \rightarrow_{\mu} F$ .

If  $x \in \bar{F}$ , it is immediate that  $x$  is a fixed point. Conversely, if  $x_0 = x$  is a fixed point, then  $x_t = x_0$ , a.e., for all  $t$ . Then, the fact  $x_t \rightarrow_{\mu} F$  implies that  $x \in \bar{F}$ . ■

We note that the fact  $x_t \rightarrow_{\mu} F$  means that the measure  $\mu_x$  associated with any limit point  $x$  of  $x_t$  is a discrete measure whose support consists of values separated by at least 1. Furthermore, it can be shown that at least one such limit point exists, because of the semi-compactness of the set of measures under the weak topology.

Theorem 5 states that  $x_t$  tends to the set  $F$ , but does not guarantee convergence to an element of this set. We make the following conjecture, which is currently unresolved.

*Conjecture 2:* Let  $(x_t)$  be a sequence of functions in  $X_L$ , evolving according to the model (5). Then, there is a function  $x^* \in F$  such that  $x_t \rightarrow_{\mu} x^*$ .

### C. Inter-Cluster Distances and Stability of Equilibria

We have found that  $x$  is a fixed point of (5) if and only if it belongs to  $\bar{F}$ , that is, with the exception of a zero-measure set, the range of  $x$  is a discrete set of values that are separated by at least one. As before, we will refer to these discrete values as clusters. In this section, we consider the stability of equilibria, and show that a condition on the inter-cluster distances similar to the one in Theorem 2 is necessary for stability. Furthermore, we show

that under a certain smoothness assumption, the system cannot converge to a fixed point that does not satisfy this condition.

In contrast to the discrete case, we can study the continuous-agent model using the classical definition of stability. We say that  $s \in F$  is *stable* if for any  $\epsilon > 0$ , there is a  $\delta > 0$  such that for any  $x_0 \in B_{\mu}(s, \delta)$ , we have  $x_t \in B_{\mu}(s, \epsilon)$  for all  $t$ . It can be shown that this notion encompasses the stability with respect to the addition of a perturbing agent used in Section II-C. More precisely, if we view the discrete-agent system as a special case of the continuum model, stability under the current definition implies stability with respect to the notion used in Section II-C. The introduction of a perturbing agent with opinion  $\tilde{x}_0$  can indeed be simulated by taking  $x_0(\alpha) = s(\alpha)$  everywhere except on an appropriate set of measure less than  $\delta$ , and  $x_0(\alpha) = \tilde{x}_0$  on this set. (However, the converse implication turns out to not hold in some pathological cases. Indeed, consider two agents separated by exactly 2. They are stable with respect to the definition of Section II-C, but not under the current definition. This is because if we introduce a small measure set of additional agents that are uniformly spread between the two original agents, we will obtain convergence to a single cluster.) Moreover, it can be verified that the notion of stability used here is equivalent to both  $\mathcal{L}_1$  and  $\mathcal{L}_2$  stability. In the sequel, and to simplify the presentation, we will neglect any zero measure sets on which  $\Delta x_t(\alpha) \neq 0$ , and will give the proof for a fixed point in  $F$ . The extension to fixed points in  $\bar{F}$  is straightforward. The proof of the following result is similar to that of its discrete counterpart, the necessary part of Theorem 2. It is omitted for space reasons, but can be found in the appendix of [7] or in [16].

*Theorem 6:* Let  $s \in F$  be a fixed point of (5), and let  $a, b$  two values taken by  $s$ . If  $s$  is stable, then

$$|b - a| \geq 1 + \frac{\min(\mu_s(a), \mu_s(b))}{\max(\mu_s(a), \mu_s(b))}. \quad (9)$$

With a little extra work, focused on the case where the distance  $|a - b|$  between the two clusters is exactly equal to 2, we can show that the strict inequality version of condition (9) is necessary for stability. We conjecture that this strict inequality version is also sufficient.

We will now proceed to show that under an additional smoothness assumption on the initial opinion function, we can never have convergence to a fixed point that violates condition (9). We start by introducing the notion of a regular opinion function. We say that a function  $x \in X_L$  is *regular* if there exist  $M \geq m > 0$  such that any interval  $J \subseteq [\inf_{\alpha} x, \sup_{\alpha} x]$  satisfies  $m|J| \leq \mu_x(J) \leq M|J|$ . Intuitively, a function is regular if the set of opinions is connected, and if the density of agents on any interval of opinions is bounded from above and from below by positive constants. (In particular, no single value is taken by a positive measure set of agents.) For example, any piecewise differentiable  $x \in X_L$  with positive upper and lower bounds on its derivative is regular.

We will show that if  $x_0$  is regular and if  $(x_t)$  converges, then  $x_t$  converges to an equilibrium satisfying the condition (9) on the minimal distance between opinions, provided that  $\sup_{\alpha} x_t - \inf_{\alpha} x_t$  remains always larger than 2. For convenience, we introduce a nonlinear update operator  $U$  on  $X_L$ , defined by  $U(x) = x - D_x^{-1} L_x x = D_x^{-1} A_x x$ , so that the recurrence (5) can be written as  $x_{t+1} = U(x_t)$ . The proof of the

following proposition is available in the appendix of [7] and in [16].

*Proposition 3:* Let  $x \in X_L$  be a regular function such that  $\sup_\alpha x - \inf_\alpha x > 2$ . Then  $U(x)$  is regular.

We note that the assumption  $\sup_\alpha x - \inf_\alpha x > 2$  in Proposition 3 is necessary for the result to hold. Indeed, if the opinion values are confined to a set  $[a, b]$ , with  $b - a = 2 - \delta < 2$ , then all agents with opinions in the set  $[a + 1 - \delta, a + 1]$  are connected with every other agent, and their next opinions will be the same, resulting in a non-regular opinion function.

As a consequence of Proposition 3, together with Theorem 5, if  $x_0$  is regular, then there are two main possibilities: i) There exists some time  $t$  at which  $\sup_\alpha x_t - \inf_\alpha x_t < 2$ . In this case, the measure  $\mu_{x_t}$  will have point masses shortly thereafter, and will eventually converge to the set of fixed points with at most two clusters. ii) Alternatively, in the ‘‘regular’’ case, we have  $\sup_\alpha x_t - \inf_\alpha x_t > 2$  for all times. Then, every  $x_t$  is regular, and convergence cannot take place in finite time. Furthermore, as we now proceed to show, convergence to a fixed point that violates the stability condition (9) is impossible. Let us note however that tight conditions for a sequence of regular functions to maintain the property  $\sup_\alpha x_t - \inf_\alpha x_t > 2$  at all times appear to be difficult to obtain.

*Theorem 7:* Let  $(x_t)$  be a sequence of functions in  $X_L$  that evolve according to (5). We assume that  $x_0$  is regular and that  $\sup_\alpha x_t - \inf_\alpha x_t > 2$  for all  $t$ . If  $(x_t)$  converges, then it converges to a function  $s \in F$  such that

$$|b - a| \geq 1 + \frac{\min(\mu_s(a), \mu_s(b))}{\max(\mu_s(a), \mu_s(b))}$$

for any two distinct values  $a, b$ , with  $\mu_s(a), \mu_s(b) > 0$ . In particular, if  $\mu_s(a) = \mu_s(b)$ , then  $|b - a| \geq 2$ .

*Proof:* Suppose that  $(x_t)$  converges to some  $s$ . It follows from Theorem 5 that  $s \in F$ , and from Proposition 3 that all  $x_t$  are regular. Suppose now that  $s$  violates the condition in the theorem, for some  $a, b$ , with  $a < b$ . Then,  $b - a < 2$ , and we must have  $\mu_s((a, b)) = 0$  because all discrete values taken by  $s$  (with positive measure) must differ by at least 1. We claim that there exists a positive length interval  $J \subseteq (a, b)$  such that  $\mu_{x_{t+1}}(J) \geq \mu_{x_t}(J)$  whenever  $x_t \in B_\mu(s, \epsilon)$ , for a sufficiently small  $\epsilon > 0$ . Since  $x_t$  converges to  $s$ , this will imply that there exists a finite time  $t^*$  after which  $\mu_{x_t}(J)$  is nondecreasing, and  $\liminf_{t \rightarrow \infty} \mu_{x_t}(J) \geq \mu_{x_{t^*}}(J) > 0$ . On the other hand, since  $\mu_s((a, b)) = 0$ ,  $\mu_{x_t}(J)$  must converge to zero. This is a contradiction and establishes the desired result.

We now establish the above claim. Let  $c = (\mu_s(a)a + \mu_s(b)b) / (\mu_s(a) + \mu_s(b))$  be the weighted average of  $a$  and  $b$ . The fact that the condition in the theorem is violated implies (c.f (3)) that  $|c - a| < 1$  and  $|c - b| < 1$ . Let  $\delta > 0$  be such that  $c - \delta + 1 > b$  and  $c + \delta - 1 < a$ , and consider the interval  $J = [c - \delta, c + \delta]$ . For any  $x \in B_\mu(s, \epsilon)$ , we have

$$\begin{aligned} \mu_x([a - \epsilon, a + \epsilon]) &\in [\mu_s(a) - \epsilon, \mu_s(a) + \epsilon], \\ \mu_x([b - \epsilon, b + \epsilon]) &\in [\mu_s(b) - \epsilon, \mu_s(b) + \epsilon], \\ \mu_x((a - 1, b + 1) \setminus ([a - \epsilon, a + \epsilon] \cup [b - \epsilon, b + \epsilon])) &\leq \epsilon \end{aligned}$$

where we have used the fact that the values taken by  $s$  are separated by at least 1. Suppose now that  $\epsilon$  is sufficiently small so that

$c - \delta + 1 > b + \epsilon$  and  $c + \delta - 1 < a - \epsilon$ . This implies that for every  $\gamma$  such that  $x(\gamma) \in J$ , we have  $(a - \epsilon, b + \epsilon) \subseteq (x(\gamma) - 1, x(\gamma) + 1)$ . If  $\epsilon$  were equal to zero, we would have  $u_x(d) = c$ . When  $\epsilon$  is small, the location of the masses at  $a$  and  $b$  moves by an  $O(\epsilon)$  amount, and an additional  $O(\epsilon)$  mass is introduced. The overall effect is easily shown to be  $O(\epsilon)$  (the detailed calculation can be found in [16]). Thus,  $|(U(x))(\gamma) - c|$  is of order  $O(\epsilon)$ . When  $\epsilon$  is chosen sufficiently small, we obtain  $c - \delta \leq (U(x))(\gamma) \leq c + \delta$ , i.e.,  $(U(x))(\gamma) \in J$  for all  $\gamma$  such that  $x(\gamma) \in J$ . This implies that  $\mu_{U(x)}(J) \geq \mu_x(J)$ , and completes the proof. ■

#### IV. RELATION BETWEEN THE DISCRETE AND THE CONTINUOUS-AGENT MODELS

We now analyze the extent to which the continuous-agent model (5) can be viewed as a limiting case of the discrete-agent model (1), when the number of agents tends to infinity. As already explained in Section III, the continuous-agent model can simulate exactly the discrete-agent model. In this section, we are interested in the converse; namely, the extent to which a discrete-agent model can describe, with arbitrarily good precision, the continuous-agent model. We will rely on the following result on the continuity of the update operator.

*Proposition 4:* Let  $x \in X_L$  be a regular function. Then, the update operator  $U$  is continuous at  $x$  with respect to the norm  $\|\cdot\|_\infty$ . More precisely, for any  $\epsilon > 0$  there exists some  $\delta > 0$  such that if  $\|y - x\|_\infty \leq \delta$  then  $\|U(y) - U(x)\|_\infty \leq \epsilon$ .

*Proof:* Consider a regular function  $x \in X_L$ , and an arbitrary  $\epsilon > 0$ . Let  $\delta$  be smaller than  $m\epsilon/25M$ , where  $m$  and  $M$  (with  $m \leq M$ ) are the bounds in the definition of regular opinion functions applied to  $x$ . We will show that if a function  $y \in X_L$  satisfies  $\|x - y\|_\infty \leq \delta$ , then  $\|U(y) - U(x)\|_\infty \leq \epsilon$ .

Fix some  $\alpha \in I$ , and let  $S_x, S_y \subseteq I$  be the set of agents connected to  $\alpha$  according to the interconnection topologies  $C_x$  and  $C_y$  defined by  $x$  and  $y$ , respectively. We let  $S_{xy} = S_x \cap S_y$ ,  $S_{x \setminus y} = S_x \setminus S_{xy}$  and  $S_{y \setminus x} = S_y \setminus S_{xy}$ . Since  $\|x - y\|_\infty \leq \delta$ , the values  $|x(\alpha) - x(\beta)|$  and  $|y(\alpha) - y(\beta)|$  differ by at most  $2\delta$ , for any  $\beta \in I$ . As a consequence, if  $\beta \in S_y$ , then  $|x(\alpha) - x(\beta)| \leq |y(\alpha) - y(\beta)| + 2\delta$ . Similarly, if  $\beta \notin S_y$ , then  $|x(\alpha) - x(\beta)| \geq |y(\alpha) - y(\beta)| - 2\delta$ . Combining these two inequalities with the definitions of  $S_{xy}$ ,  $S_{x \setminus y}$ , and  $S_{y \setminus x}$ , we obtain

$$\begin{aligned} [x(\alpha) - 1 + 2\delta, x(\alpha) + 1 - 2\delta] &\subseteq x(S_{xy}) \subseteq [x(\alpha) - 1, x(\alpha) + 1], \\ x(S_{x \setminus y}) &\subseteq [x(\alpha) - 1, x(\alpha) - 1 + 2\delta] \cup [x(\alpha) + 1 - 2\delta, x(\alpha) + 1], \\ x(S_{y \setminus x}) &\subseteq [x(\alpha) - 1 - 2\delta, x(\alpha) - 1] \cup [x(\alpha) + 1, x(\alpha) + 1 + 2\delta]. \end{aligned}$$

Since  $x$  is regular, we have  $|S_{xy}| \geq m(2 - 4\delta) \geq m$  and  $|S_{x \setminus y}|, |S_{y \setminus x}| \leq M4\delta$ . Let now  $\bar{x}_{xy}$  and  $\bar{x}_{x \setminus y}$  be the average value of  $x$  on  $S_{xy}$  and  $S_{x \setminus y}$ , respectively. Similarly, let  $\bar{y}_{xy}$ , and  $\bar{y}_{y \setminus x}$  be the average value of  $y$  on  $S_{xy}$  and  $S_{y \setminus x}$ . Since  $\|x - y\|_\infty \leq \delta$ ,  $\bar{x}_{xy}$  and  $\bar{y}_{xy}$  differ by at most  $\delta$ . It follows from the definition of the model (5) that:

$$\begin{aligned} (U(x))(\alpha) &= \bar{x}_{xy} + \frac{|S_{x \setminus y}|}{|S_{xy}| + |S_{x \setminus y}|} (\bar{x}_{x \setminus y} - \bar{x}_{xy}), \\ (U(y))(\alpha) &= \bar{y}_{xy} + \frac{|S_{y \setminus x}|}{|S_{xy}| + |S_{y \setminus x}|} (\bar{y}_{y \setminus x} - \bar{y}_{xy}). \end{aligned}$$

It can be seen that  $|\bar{x}_{x\setminus y} - \bar{x}_{xy}| \leq 3$  and  $|\bar{y}_{y\setminus x} - \bar{y}_{xy}| \leq 3$ , from which we obtain that  $|(U(y))(\alpha) - (U(x))(\alpha)|$  is upper

$$|\bar{x}_{xy} - \bar{y}_{xy}| + 3 \frac{|S_{y\setminus x}|}{|S_{xy}|} + 3 \frac{|S_{x\setminus y}|}{|S_{xy}|} \leq \delta + 6 \frac{4M\delta}{m} \leq \epsilon,$$

where we have used the fact that  $|\bar{x}_{xy} - \bar{y}_{xy}| \leq \delta$ . Since the above is true for any  $\alpha \in I$ , we conclude that  $\|U(y) - U(x)\|_\infty \leq \epsilon$ . ■

Let  $U^t : X_L \rightarrow X_L$  be the composition of the update operator, defined by  $U^t(x) = U(U^{t-1}(x))$ , so that  $U^t(x_0) = x_t$ . Proposition 4 is readily extended to a continuity result for  $U^t$ .

*Corollary 1:* Let  $x_0 \in X_L$  be a regular function such that  $\sup_\alpha U^t(x) - \inf_\alpha U^t(x) > 2$  for every  $t \geq 0$ . Then for any finite  $t$ ,  $U^t$  is continuous at  $x$  with respect to the norm  $\|\cdot\|_\infty$ .

*Proof:* Since  $x$  is regular and since  $\sup_\alpha U^t(x) - \inf_\alpha U^t(x) > 2$  for all  $t$ , Proposition 3 implies that all  $U^t(x)$  are regular. Proposition 4 then implies that for all  $t$ ,  $U$  is continuous at  $U^t(x)$ , and therefore the composition  $U^t$  is continuous at  $x$ . ■

Corollary 1 allows us to prove that, in the regular case, and for any given finite time horizon, the continuous-agent model is the limit of the discrete-agent model, as the number of agents grows. To this effect, for any given partition of  $I = [0, 1]$  into  $n$  disjoint sets  $J_1, \dots, J_n$ , we define an operator  $\mathcal{G} : \mathfrak{R}^n \rightarrow X$  that translates the opinions in an  $n$ -agent system to an opinion function in the continuous-agent model. More precisely, for a vector  $\hat{x} \in \mathfrak{R}^n$  and any  $\alpha \in J_i$ , we let  $(\mathcal{G}\hat{x})(\alpha)$  be equal to the  $i$ th component of  $\hat{x}$ .

*Theorem 8:* Let  $x_0 \in X_L$  be a regular function and assume that  $\sup_\alpha x_t - \inf_\alpha x_t > 2$  for  $t \leq t^*$ . Then, the sequence  $(x_t)$ ,  $t = 1, \dots, t^*$ , can be approximated arbitrarily well by a sequence  $(\hat{x}_t)$  of opinion vectors evolving according to (1), in the following sense. For any  $\epsilon > 0$ , there exists some  $n$ , a partition of  $I$  into  $n$  disjoint sets  $J_1, \dots, J_n$ , and a vector  $\hat{x}_0 \in [0, L]^n$  such that the sequence of vectors  $\hat{x}_t$  generated by the discrete-agent model (1), starting from  $\hat{x}_0$ , satisfies  $\|x_t - \mathcal{G}\hat{x}_t\|_\infty \leq \epsilon$ , for  $t = 1, \dots, t^*$ .

*Proof:* Fix  $\epsilon > 0$ . Since all  $U^t$  are continuous at  $x_0$ , there is some  $\delta > 0$  such that if  $\|y - x_0\|_\infty \leq \delta$ , then  $\|U^t(y) - x_t\|_\infty \leq \epsilon$ , for  $t \leq t^*$ . Since  $x_0$  is regular, we can divide  $I$  into subsets  $J_1, J_2, \dots, J_n$ , so that  $|J_i| = 1/n$  for all  $i$ , and  $|x_0(\alpha) - x_0(\beta)| \leq \delta$  for all  $\alpha, \beta$  in the same set  $J_i$ . (This is done by letting  $c_i$  be such that  $\mu_{x_0}([0, c_i]) = i/n$ , and defining  $J_i = \{\alpha : c_{i-1} < x_0(\alpha) \leq c_i\}$ , where  $n$  is sufficiently large.) We define  $\hat{x}_0 \in [0, L]^n$  by letting its  $i$ th component be equal to  $c_i$ . We then have  $\|x_0 - \mathcal{G}\hat{x}_0\|_\infty \leq \delta$ . This implies that  $\|x_t - U^t(\mathcal{G}\hat{x}_0)\|_\infty \leq \epsilon$ , for  $t \leq t^*$ . Since the continuous-agent model, initialized with a discrete distribution, simulates the discrete-agent model, we have  $U^t(\mathcal{G}\hat{x}_0) = \mathcal{G}\hat{x}_t$ , and the desired result follows. ■

Theorem 8 supports the intuition that for large values of  $n$ , the continuous-agent model behaves approximatively as the discrete-agent model, over any finite horizon. In view of Theorem 6, this suggests that the discrete-agent system should always converge to a stable equilibrium (in the sense defined in Section II) when  $n$  is sufficiently large, as stated in Conjecture 1, and observed in many examples (see, e.g., Fig. 6). Indeed, Theorem 6 states that under the regularity assumption, the continuum system cannot converge to an equilibrium that does not

satisfy condition (9) on the inter-cluster distances. However, this argument does not translate to a proof of the conjecture because the approximation property in Theorem 8 only holds over a finite time horizon, and does not necessarily provide information on the limiting behavior.

## V. CONCLUSION

We have analyzed the model of opinion dynamics (1) introduced by Krause, from several angles. Our motivation was to provide an analysis of a simple multi-agent system with an endogenously changing interconnection topology while taking explicitly advantage of the topology dynamics, something that is rarely done in the related literature.

We focused our attention on an intriguing phenomenon, the fact that equilibrium inter-cluster distances are usually significantly larger than 1, and typically close to two. We proposed an explanation of this phenomenon based on a notion of stability with respect to the addition of a perturbing agent. We showed that such stability translates to a certain lower bound on the inter-cluster distances, with the bound equal to two when the clusters have identical weights. We also discussed the conjecture that when the number of agents is sufficiently large, the system converges to a stable equilibrium for “most” initial conditions.

To avoid granularity problems linked with the presence or absence of an agent in a particular region, we introduced a new opinion dynamics model that allows for a continuum of agents. For this model we proved that under some regularity assumptions, there is always a finite density of agents between any two clusters during the convergence process. As a result, we could prove that such systems never converge to an unstable equilibrium. We also proved that the continuous-agent model is indeed the limit of a discrete model, over any given finite time horizon, as the number of agents grows to infinity. These results provide some additional support for the conjectured, but not yet established, generic convergence to stable equilibria.

We originally introduced the continuous-agent model as a tool for the study of the discrete-agent model, but it is also of independent interest and raises some challenging open questions. An important one is the question of whether the continuous-agent model is always guaranteed to converge. (We only succeeded in establishing convergence to the set of fixed points, not to a single fixed point.)

Finally, the study of the continuous-agent model suggests some broader questions. In the same way that the convergence of the discrete-agent model can be viewed as a special case of convergence of inhomogeneous products of stochastic matrices, it may be fruitful to view the convergence of the continuous-agent model as a special case of convergence of inhomogeneous compositions of stochastic operators, and to develop results for the latter problem.

The model (1) can of course be extended to higher dimensional spaces, as is often done in the opinion dynamics literature (see [25] for a survey). Numerical experiments again show the emergence of clusters that are separated by distances significantly larger than 1. The notion of stability with respect to the addition of an agent can also be extended to higher dimensions. However, stability conditions become more complicated,

and in particular cannot be expressed as a conjunction of independent conditions, one for each pair of clusters. For example, it turns out that adding a cluster to an unstable equilibrium may render it stable [16]. In addition, a formal analysis appears difficult because in  $\mathfrak{R}^n$ , with  $n > 1$ , the support of the opinion distribution can be connected without being convex, and convexity is not necessarily preserved by our systems. For this reason, even under “regularity” assumptions, the presence of perturbing agents between clusters is not guaranteed.

## REFERENCES

- [1] E. Ben-Naim, “Rise and fall of political parties,” *Europhys. Lett.*, vol. 69, no. 5, pp. 671–676, 2005.
- [2] E. Ben-Naim, P. L. Krapivsky, and S. Redner, “Bifurcations and patterns in compromise processes,” *Physica D*, vol. 183, no. 3, pp. 190–204, 2003.
- [3] E. Ben-Naim, P. L. Krapivsky, F. Vasquez, and S. Redner, “Unity and discord in opinion dynamics,” *Physica A*, vol. 330, pp. 99–106, 2003.
- [4] D. P. Bertsekas and J. N. Tsitsiklis, *Parallel and Distributed Computation*. Englewood Cliffs, NJ: Prentice-Hall, 1989.
- [5] V. D. Blondel, J. M. Hendrickx, A. Olshevsky, and J. N. Tsitsiklis, “Convergence in multiagent coordination, consensus, and flocking,” in *Proc. 44th IEEE Conf. Decision Control (CDC'2005)*, Seville, Spain, Dec. 2005, pp. 2996–3000.
- [6] V. D. Blondel, J. M. Hendrickx, and J. N. Tsitsiklis, “On the 2R conjecture for multi-agent systems,” in *Proc. Eur. Control Conf. (ECC'2007)*, Kos, Greece, Jul. 2007, pp. 874–881.
- [7] V. D. Blondel, J. M. Hendrickx, and J. N. Tsitsiklis, “On Krause’s Consensus Multi-Agent Model With State-Dependent Connectivity (Extended Version),” *Comput. Sci. Multiagent Syst.*, 2009 [Online]. Available: <http://arxiv.org/abs/0807.2028>
- [8] F. Cucker and S. Smale, “Emergent behavior in flocks,” *IEEE Trans. Autom. Control*, vol. 52, no. 5, pp. 852–862, May 2007.
- [9] F. Cucker and S. Smale, “On the mathematics of emergence,” *Japanese J. Math.*, vol. 2, pp. 119–227, Mar. 2007.
- [10] G. Deffuant, D. Neau, F. Amblard, and G. Weisbuch, “Mixing beliefs among interacting agents,” *Adv. Complex Syst.*, vol. 3, pp. 87–98, 2000.
- [11] M. H. DeGroot, “Reaching a consensus,” *J. Amer. Stat. Assoc.*, vol. 69, pp. 118–121, 1974.
- [12] J. C. Dittmer, “Consensus formation under bounded confidence,” *Nonlin. Anal.*, no. 47, pp. 4615–4621, 2001.
- [13] S. Fortunato, V. Latora, A. Pluchino, and R. Rapisarda, “Vector opinion dynamics in a bounded confidence consensus model,” *Int. J. Modern Phys. C*, vol. 16, pp. 1535–1551, Oct. 2005.
- [14] R. Hegselmann, “Opinion dynamics: Insights by radically simplifying models,” in *Laws and Models in Science*, D. Gillies, Ed. London, U.K.: Kings College, 2004, pp. 1–29.
- [15] R. Hegselmann and U. Krause, “Opinion dynamics and bounded confidence models, analysis, and simulations,” *J. Artif. Societies Social Simul.*, vol. 5, no. 3, 2002.
- [16] J. M. Hendrickx, “Graphs and Networks for the Analysis of Autonomous Agent Systems” Ph.D. dissertation, Université catholique de Louvain, Louvain, Belgium, 2008.
- [17] J. M. Hendrickx and V. D. Blondel, “Convergence of different linear and non-linear Vicsek models,” in *Proc. 17th Int. Symp. Math. Theory Networks Syst. (MTNS'06)*, Kyoto, Japan, Jul. 2006, pp. 1229–1240.
- [18] A. Jadbabaie, J. Lin, and A. S. Morse, “Coordination of groups of mobile autonomous agents using nearest neighbor rules,” *IEEE Trans. Autom. Control*, vol. 48, no. 6, pp. 988–1001, Jun. 2003.
- [19] E. W. Justh and P. S. Krishnaprasad, “Equilibria and steering laws for planar formations,” *Syst. Control Lett.*, vol. 52, no. 1, pp. 25–38, 2004.
- [20] U. Krause, “Soziale dynamiken mit vielen interakteuren. Eine problemskizze,” in *Proc. Modellierung Simul. von Dynamiken mit vielen interagierenden Akteuren*, 1997, pp. 37–51.
- [21] U. Krause, “A discrete nonlinear and non-autonomous model of consensus formation,” in *Proc. Commun. Difference Equations*, 2000, pp. 227–236.
- [22] J. Lin, A. S. Morse, and B. D. O. Anderson, “The multi-agent rendezvous problem,” in *Proc. 42th IEEE Conf. Decision Control (CDC'03)*, Honolulu, HI, Dec. 2003, pp. 1508–1513.
- [23] J. Lorenz, “A stabilization theorem for continuous opinion dynamics,” *Physica A*, vol. 355, no. 1, pp. 217–223, 2005.
- [24] J. Lorenz, “Consensus strikes back in the Hegselmann-Krause model of continuous opinion dynamics under bounded confidence,” *J. Artif. Societies Social Simul.*, vol. 9, no. 1, 2006.
- [25] J. Lorenz, “Continuous opinion dynamics under bounded confidence: A survey,” *Int. J. Modern Phys. C*, vol. 18, no. 12, pp. 1819–1838, 2007.
- [26] L. Moreau, “Stability of multiagent systems with time-dependent communication links,” *IEEE Trans. Autom. Control*, vol. 50, no. 2, pp. 169–182, Feb. 2005.
- [27] R. Olfati-Saber, J. A. Fax, and R. M. Murray, “Consensus and cooperation in networked multi-agent systems,” *Proc. IEEE*, vol. 95, no. 1, pp. 215–233, Jan. 2007.
- [28] W. Ren, R. W. Beard, and E. M. Atkins, *IEEE Control Syst. Mag.*, vol. 27, no. 2, pp. 71–82, Apr. 2007.
- [29] J. N. Tsitsiklis, “Problems in Decentralized Decision Making and Computation” Ph.D. dissertation, Dept. Elect. Eng. Comput. Sci., Massachusetts Inst. Technol., Cambridge, 1984.
- [30] D. Urbig, “Attitude dynamics with limited verbalisation capabilities,” *J. Artif. Societies Social Simul.*, vol. 6, no. 1, 2003.
- [31] T. Vicsek, A. Czirok, I. Ben Jacob, I. Cohen, and O. Schochet, “Novel type of phase transitions in a system of self-driven particles,” *Phys. Rev. Lett.*, vol. 75, pp. 1226–1229, 1995.



**Vincent D. Blondel** received the M.Sc. degree in mathematics from Imperial College, London, U.K., in 1990 and the Ph.D. degree in applied mathematics from the Université catholique de Louvain, Louvain-la-Neuve, Belgium, in 1992.

He was a Visiting Researcher at the Royal Institute of Technology, Stockholm, Sweden, and at the Institut National de Recherche en Informatique et en Automatique (INRIA), Rocquencourt, France. From 2005 to 2006, he was an Invited Professor and a Fulbright Scholar at the Massachusetts Institute of Technology, Cambridge. He is currently a Professor and Department Head at the Université catholique de Louvain.

Dr. Blondel received the Prize Wetrems of the Belgian Royal Academy of Science, the Society for Industrial and Applied Mathematics (SIAM) Prize on Control and Systems Theory, and the Ruberti Prize in Systems and Control of the IEEE in 2006.



**Julien M. Hendrickx** received the M.Eng. degree in applied mathematics and the Ph.D. degree in mathematical engineering from the Université catholique de Louvain, Louvain, Belgium, in 2004 and 2008, respectively.

He has been a Visiting Researcher at the University of Illinois at Urbana Champaign, from 2003 to 2004, at the National ICT Australia in 2005 and 2006, and at the Massachusetts Institute of Technology (MIT), Cambridge, in 2006 and 2008. He is currently a Postdoctoral Fellow at the Laboratory for Information and

Decision Systems, MIT, and holds postdoctoral fellowships of the F.R.S.-FNRS (Fund for Scientific Research) and the Belgian American Education Foundation.

Dr. Hendrickx received the 2008 EECI award for the best Ph.D. thesis in Europe in the field of embedded and networked control.



**John N. Tsitsiklis** (F'99) received the B.S. degree in mathematics and the B.S., M.S., and Ph.D. degrees in electrical engineering from the Massachusetts Institute of Technology (MIT), Cambridge, in 1980, 1980, 1981, and 1984, respectively.

He is currently a Clarence J. Lebel Professor with the Department of Electrical Engineering, MIT. He has served as a Codirector of the MIT Operations Research Center from 2002 to 2005, and in the National Council on Research and Technology in Greece (2005 to 2007). His research interests are in

systems, optimization, communications, control, and operations research. He has coauthored four books and more than 100 journal papers in these areas.

Dr. Tsitsiklis received the Outstanding Paper Award from the IEEE Control Systems Society (1986), the M.I.T. Edgerton Faculty Achievement Award (1989), the Bodossakis Foundation Prize (1995), and the INFORMS/CSTS Prize (1997). He is a member of the National Academy of Engineering. Finally, in 2008, he was conferred the title of Doctor honoris causa, from the Université catholique de Louvain