

Distributed Optimization via Alternating Direction Method of Multipliers

Stephen Boyd, Neal Parikh, Eric Chu, Borja Peleato

Stanford University

ITMANET, Stanford, January 2011

Outline

- precursors
 - dual decomposition
 - method of multipliers
- alternating direction method of multipliers
- applications/examples
- conclusions/big picture

Dual problem

- convex equality constrained optimization problem

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax = b \end{array}$$

- Lagrangian: $L(x, y) = f(x) + y^T(Ax - b)$
- dual function: $g(y) = \inf_x L(x, y)$
- dual problem: maximize $g(y)$
- recover $x^* = \operatorname{argmin}_x L(x, y^*)$

Dual ascent

- gradient method for dual problem: $y^{k+1} = y^k + \alpha^k \nabla g(y^k)$
- $\nabla g(y^k) = A\tilde{x} - b$, where $\tilde{x} = \operatorname{argmin}_x L(x, y^k)$
- dual ascent method is

$$x^{k+1} := \operatorname{argmin}_x L(x, y^k) \quad // \textit{x-minimization}$$

$$y^{k+1} := y^k + \alpha^k (Ax^{k+1} - b) \quad // \textit{dual update}$$

- works, with lots of strong assumptions

Dual decomposition

- suppose f is separable:

$$f(x) = f_1(x_1) + \cdots + f_N(x_N), \quad x = (x_1, \dots, x_N)$$

- then L is separable in x : $L(x, y) = L_1(x_1, y) + \cdots + L_N(x_N, y) - y^T b$,

$$L_i(x_i, y) = f_i(x_i) + y^T A_i x_i$$

- x -minimization in dual ascent splits into N separate minimizations

$$x_i^{k+1} := \operatorname{argmin}_{x_i} L_i(x_i, y^k)$$

which can be carried out in parallel

Dual decomposition

- dual decomposition (*Everett, Dantzig, Wolfe, Benders 1960–65*)

$$x_i^{k+1} := \operatorname{argmin}_{x_i} L_i(x_i, y^k), \quad i = 1, \dots, N$$

$$y^{k+1} := y^k + \alpha^k (\sum_{i=1}^N A_i x_i^{k+1} - b)$$

- scatter y^k ; update x_i in parallel; gather $A_i x_i^{k+1}$
- solve a large problem
 - by iteratively solving subproblems (in parallel)
 - dual variable update provides coordination
- works, with lots of assumptions; often slow

Method of multipliers

- a method to robustify dual ascent
- use **augmented Lagrangian** (*Hestenes, Powell 1969*), $\rho > 0$

$$L_\rho(x, y) = f(x) + y^T(Ax - b) + (\rho/2)\|Ax - b\|_2^2$$

- method of multipliers (*Hestenes, Powell; analysis in Bertsekas 1982*)

$$\begin{aligned}x^{k+1} &:= \operatorname{argmin}_x L_\rho(x, y^k) \\ y^{k+1} &:= y^k + \rho(Ax^{k+1} - b)\end{aligned}$$

(note specific dual update step length ρ)

Method of multipliers

- good news: converges under much more relaxed conditions (f can be nondifferentiable, take on value $+\infty$, . . .)
- bad news: quadratic penalty destroys splitting of the x -update, so can't do decomposition

Alternating direction method of multipliers

- a method
 - with good robustness of method of multipliers
 - which can support decomposition

“robust dual decomposition” or “decomposable method of multipliers”
- proposed by Gabay, Mercier, Glowinski, Marrocco in 1976

Alternating direction method of multipliers

- ADMM problem form (with f, g convex)

$$\begin{aligned} & \text{minimize} && f(x) + g(z) \\ & \text{subject to} && Ax + Bz = c \end{aligned}$$

– two sets of variables, with separable objective

- $L_\rho(x, z, y) = f(x) + g(z) + y^T(Ax + Bz - c) + (\rho/2)\|Ax + Bz - c\|_2^2$
- ADMM:

$$x^{k+1} := \operatorname{argmin}_x L_\rho(x, z^k, y^k) \quad // \text{ } x\text{-minimization}$$

$$z^{k+1} := \operatorname{argmin}_z L_\rho(x^{k+1}, z, y^k) \quad // \text{ } z\text{-minimization}$$

$$y^{k+1} := y^k + \rho(Ax^{k+1} + Bz^{k+1} - c) \quad // \text{ dual update}$$

Alternating direction method of multipliers

- if we minimized over x and z jointly, reduces to method of multipliers
- instead, we do one pass of a Gauss-Seidel method
- we get splitting since we minimize over x with z fixed, and vice versa

Convergence

- assume (very little!)
 - f, g convex, closed, proper
 - L_0 has a saddle point
- then ADMM converges:
 - iterates approach feasibility: $Ax^k + Bz^k - c \rightarrow 0$
 - objective approaches optimal value: $f(x^k) + g(z^k) \rightarrow p^*$

Related algorithms

- operator splitting methods
(*Douglas, Peaceman, Rachford, Lions, Mercier, . . . 1950s, 1979*)
- proximal point algorithm (*Rockafellar 1976*)
- Dykstra's alternating projections algorithm (1983)
- Spingarn's method of partial inverses (1985)
- Rockafellar-Wets progressive hedging (1991)
- proximal methods (*Rockafellar, many others, 1976–present*)
- Bregman iterative methods (2008–present)
- most of these are special cases of the proximal point algorithm

Consensus optimization

- want to solve problem with N objective terms

$$\text{minimize } \sum_{i=1}^N f_i(x)$$

- *e.g.*, f_i is the loss function for i th block of training data

- ADMM form:

$$\begin{aligned} &\text{minimize } \sum_{i=1}^N f_i(x_i) \\ &\text{subject to } x_i - z = 0 \end{aligned}$$

- x_i are *local variables*
- z is the *global variable*
- $x_i - z = 0$ are *consistency* or *consensus* constraints

Consensus optimization via ADMM

- $L_\rho(x, z, y) = \sum_{i=1}^N (f_i(x_i) + y_i^T(x_i - z) + (\rho/2)\|x_i - z\|_2^2)$
- ADMM:

$$x_i^{k+1} := \operatorname{argmin}_{x_i} (f_i(x_i) + y_i^{kT}(x_i - z^k) + (\rho/2)\|x_i - z^k\|_2^2)$$

$$z^{k+1} := \frac{1}{N} \sum_{i=1}^N (x_i^{k+1} + (1/\rho)y_i^k)$$

$$y_i^{k+1} := y_i^k + \rho(x_i^{k+1} - z^{k+1})$$

Consensus optimization via ADMM

- using $\sum_{i=1}^N y_i^k = 0$, algorithm simplifies to

$$x_i^{k+1} := \operatorname{argmin}_{x_i} (f_i(x_i) + y_i^{kT} (x_i - \bar{x}^k) + (\rho/2) \|x_i - \bar{x}^k\|_2^2)$$

$$y_i^{k+1} := y_i^k + \rho(x_i^{k+1} - \bar{x}^{k+1})$$

where $\bar{x}^k = (1/N) \sum_{i=1}^N x_i^k$

- in each iteration
 - gather x_i^k and average to get \bar{x}^k
 - scatter the average \bar{x}^k to processors
 - update y_i^k locally (in each processor, in parallel)
 - update x_i locally

Statistical interpretation

- f_i is negative log-likelihood for parameter x given i th data block
- x_i^{k+1} is MAP estimate under prior $\mathcal{N}(\bar{x}^k + (1/\rho)y_i^k, \rho I)$
- prior mean is previous iteration's consensus shifted by 'price' of processor i disagreeing with previous consensus
- processors only need to support a Gaussian MAP method
 - type or number of data in each block not relevant
 - consensus protocol yields global maximum-likelihood estimate


Consensus classification

- data (examples) (a_i, b_i) , $i = 1, \dots, N$, $a_i \in \mathbf{R}^n$, $b_i \in \{-1, +1\}$
- linear classifier $\text{sign}(a^T w + v)$, with weight w , offset v
- margin for i th example is $b_i(a_i^T w + v)$; want margin to be positive
- loss for i th example is $l(b_i(a_i^T w + v))$
 - l is loss function (hinge, logistic, probit, exponential, . . .)
- choose w, v to minimize $\frac{1}{N} \sum_{i=1}^N l(b_i(a_i^T w + v)) + r(w)$
 - $r(w)$ is regularization term (ℓ_2, ℓ_1, \dots)
- split data and use ADMM consensus to solve


Consensus SVM example

- hinge loss $l(u) = (1 - u)_+$ with ℓ_2 regularization
- baby problem with $n = 2$, $N = 400$ to illustrate
- examples split into 20 groups, in worst possible way:
each group contains only positive or negative examples


Iteration 1



Iteration 5



Iteration 40



Reference

Distributed Optimization and Statistical Learning via the Alternating Direction Method of Multipliers (Boyd, Parikh, Chu, Peleato, Eckstein)

available at Boyd web site