

# Bayesian Interpretations of Regularization

Charlie Frogner

9.520 Class 17

April 6, 2011

Regularized least squares maps  $\{(x_i, y_i)\}_{i=1}^n$  to a function that minimizes the regularized loss:

$$f_S = \arg \min_{f \in \mathcal{H}} \frac{1}{2} \sum_{i=1}^n (y_i - f(x_i))^2 + \frac{\lambda}{2} \|f\|_{\mathcal{H}}^2$$

*Can we interpret RLS from a probabilistic point of view?*

# Some notation

- Training set:  $S = \{(\mathbf{x}_1, y_1), \dots, (\mathbf{x}_n, y_n)\}$ .
- Inputs:  $\mathbf{X} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ .
- Labels:  $\mathbf{Y} = \{y_1, \dots, y_n\}$ .
- Parameters:  $\theta \in \mathbb{R}^p$ .
- $p(\mathbf{Y}|\mathbf{X}, \theta)$  is the joint distribution over labels  $\mathbf{Y}$  given inputs  $\mathbf{X}$  and the parameters.

# Where do probabilities show up?

$$\frac{1}{2} \sum_{i=1}^n V(y_i, f(x_i)) + \frac{\lambda}{2} \|f\|_{\mathcal{H}}^2$$

becomes

$$p(\mathbf{Y}|f, \mathbf{X}) \cdot p(f)$$

- **Likelihood**, a.k.a. **noise model**:  $p(\mathbf{Y}|f, \mathbf{X})$ .
  - Gaussian:  $y_i \sim \mathcal{N}(f^*(x_i), \sigma_i^2)$
  - Poisson:  $y_i \sim \text{Pois}(f^*(x_i))$
- **Prior**:  $p(f)$ .

# Where do probabilities show up?

$$\frac{1}{2} \sum_{i=1}^n V(y_i, f(\mathbf{x}_i)) + \frac{\lambda}{2} \|f\|_{\mathcal{H}}^2$$

becomes

$$p(\mathbf{Y}|f, \mathbf{X}) \cdot p(f)$$

- **Likelihood**, a.k.a. **noise model**:  $p(\mathbf{Y}|f, \mathbf{X})$ .
  - Gaussian:  $y_i \sim \mathcal{N}(f^*(\mathbf{x}_i), \sigma_i^2)$
  - Poisson:  $y_i \sim \text{Pois}(f^*(\mathbf{x}_i))$
- **Prior**:  $p(f)$ .

The estimation problem:

- Given data  $\{(x_i, y_i)\}_{i=1}^N$  and model  $p(\mathbf{Y}|f, \mathbf{X}), p(f)$ .
- Find a good  $f$  to explain data.

- Maximum likelihood estimation for ERM
- MAP estimation for linear RLS
- MAP estimation for kernel RLS
- Transductive model
- Infinite dimensions get more complicated

# Maximum likelihood estimation

- Given data  $\{(x_i, y_i)\}_{i=1}^N$  and model  $p(\mathbf{Y}|f, \mathbf{X}), p(f)$ .
- A good  $f$  is one that maximizes  $p(\mathbf{Y}|f, \mathbf{X})$ .



# Maximum likelihood and least squares

For least squares, noise model is:

$$y_i | f, \mathbf{x}_i \sim \mathcal{N}(f(\mathbf{x}_i), \sigma^2)$$

a.k.a.

$$\mathbf{Y} | f, \mathbf{X} \sim \mathcal{N}(f(\mathbf{X}), \sigma^2 I)$$

So

$$p(\mathbf{Y} | f, \mathbf{X}) = \frac{1}{(2\pi\sigma^2)^{N/2}} \exp \left\{ - \sum_{i=1}^N \frac{1}{\sigma^2} (y_i - f(x_i))^2 \right\}$$

# Maximum likelihood and least squares

For least squares, noise model is:

$$y_i | f, \mathbf{x}_i \sim \mathcal{N}(f(\mathbf{x}_i), \sigma^2)$$

a.k.a.

$$\mathbf{Y} | f, \mathbf{X} \sim \mathcal{N}(f(\mathbf{X}), \sigma^2 I)$$

So

$$p(\mathbf{Y} | f, \mathbf{X}) = \frac{1}{(2\pi\sigma^2)^{N/2}} \exp \left\{ - \sum_{i=1}^N \frac{1}{\sigma^2} (y_i - f(\mathbf{x}_i))^2 \right\}$$

# Maximum likelihood and least squares

Maximum likelihood: maximize

$$p(\mathbf{Y}|f, \mathbf{X}) = \frac{1}{(2\pi\sigma^2)^{N/2}} \exp \left\{ - \sum_{i=1}^N \frac{1}{\sigma^2} (y_i - f(\mathbf{x}_i))^2 \right\}$$

Empirical risk minimization: minimize

$$\sum_{i=1}^N (y_i - f(\mathbf{x}_i))^2$$

$$\sum_{i=1}^N (y_i - f(x_i))^2$$

$$e^{-\sum_{i=1}^N \frac{1}{\sigma^2} (y_i - f(x_i))^2}$$

# What about regularization?

RLS:

$$\arg \min_f \frac{1}{2} \sum_{i=1}^n (y_i - f(x_i))^2 + \frac{\lambda}{2} \|f\|_{\mathcal{H}}^2$$

Is there a model of  $\mathbf{Y}$  and  $f$  that yields RLS?

Yes.

$$e^{-\frac{1}{2\sigma_\varepsilon^2} \left( \sum_{i=1}^n (y_i - f(x_i))^2 \right) - \frac{\lambda}{2} \|f\|_{\mathcal{H}}^2}$$

$$p(\mathbf{Y}|f, \mathbf{X}) \cdot p(f)$$

# What about regularization?

RLS:

$$\arg \min_f \frac{1}{2} \sum_{i=1}^n (y_i - f(x_i))^2 + \frac{\lambda}{2} \|f\|_{\mathcal{H}}^2$$

Is there a model of  $\mathbf{Y}$  and  $f$  that yields RLS?

Yes.

$$e^{-\frac{1}{2\sigma_\varepsilon^2} \left( \sum_{i=1}^n (y_i - f(x_i))^2 \right) - \frac{\lambda}{2} \|f\|_{\mathcal{H}}^2}$$

$$p(\mathbf{Y}|f, \mathbf{X}) \cdot p(f)$$

# What about regularization?

RLS:

$$\arg \min_f \frac{1}{2} \sum_{i=1}^n (y_i - f(x_i))^2 + \frac{\lambda}{2} \|f\|_{\mathcal{H}}^2$$

Is there a model of  $\mathbf{Y}$  and  $f$  that yields RLS?

Yes.

$$e^{-\frac{1}{2\sigma_\varepsilon^2} \left( \sum_{i=1}^n (y_i - f(x_i))^2 \right)} \cdot e^{-\frac{\lambda}{2} \|f\|_{\mathcal{H}}^2}$$

$$p(\mathbf{Y}|f, \mathbf{X}) \cdot p(f)$$



# What about regularization?

RLS:

$$\arg \min_f \frac{1}{2} \sum_{i=1}^n (y_i - f(x_i))^2 + \frac{\lambda}{2} \|f\|_{\mathcal{H}}^2$$

Is there a model of  $\mathbf{Y}$  and  $f$  that yields RLS?

Yes.

$$e^{-\frac{1}{2\sigma^2\varepsilon} \left( \sum_{i=1}^n (y_i - f(x_i))^2 \right)} \cdot e^{-\frac{\lambda}{2} \|f\|_{\mathcal{H}}^2}$$

$$p(\mathbf{Y}|f, \mathbf{X}) \cdot p(f)$$

# Posterior function estimates

- Given data  $\{(x_i, y_i)\}_{i=1}^N$  and model  $p(\mathbf{Y}|f, \mathbf{X}), p(f)$ .
- Find a good  $f$  to explain data.

(If we can get  $p(f|\mathbf{Y}, \mathbf{X})$ )

*Bayes least squares estimate:*

$$\hat{f}_{BLS} = \mathbb{E}_{(f|\mathbf{X}, \mathbf{Y})}[f]$$

i.e. the mean of the posterior.

*MAP estimate:*

$$\hat{f}_{MAP}(\mathbf{Y}|\mathbf{X}) = \arg \max_f p(f|\mathbf{X}, \mathbf{Y})$$

i.e. a mode of the posterior.

# Posterior function estimates

- Given data  $\{(x_i, y_i)\}_{i=1}^N$  and model  $p(\mathbf{Y}|f, \mathbf{X}), p(f)$ .
- Find a good  $f$  to explain data.

(If we can get  $p(f|\mathbf{Y}, \mathbf{X})$ )

*Bayes least squares estimate:*

$$\hat{f}_{BLS} = \mathbb{E}_{(f|\mathbf{X}, \mathbf{Y})}[f]$$

i.e. the mean of the posterior.

*MAP estimate:*

$$\hat{f}_{MAP}(\mathbf{Y}|\mathbf{X}) = \arg \max_f p(f|\mathbf{X}, \mathbf{Y})$$

i.e. a mode of the posterior.

# Posterior function estimates

- Given data  $\{(x_i, y_i)\}_{i=1}^N$  and model  $p(\mathbf{Y}|f, \mathbf{X}), p(f)$ .
- Find a good  $f$  to explain data.

(If we can get  $p(f|\mathbf{Y}, \mathbf{X})$ )

*Bayes least squares estimate:*

$$\hat{f}_{BLS} = \mathbb{E}_{(f|\mathbf{X}, \mathbf{Y})}[f]$$

i.e. the mean of the posterior.

*MAP estimate:*

$$\hat{f}_{MAP}(\mathbf{Y}|\mathbf{X}) = \arg \max_f p(f|\mathbf{X}, \mathbf{Y})$$

i.e. a mode of the posterior.

# A posterior on functions?

How to find  $p(f|\mathbf{Y}, \mathbf{X})$ ?

*Bayes' rule:*

$$\begin{aligned} p(f|\mathbf{X}, \mathbf{Y}) &= \frac{p(\mathbf{Y}|\mathbf{X}, f) \cdot p(f)}{p(\mathbf{Y}|\mathbf{X})} \\ &= \frac{p(\mathbf{Y}|\mathbf{X}, f) \cdot p(f)}{\int p(\mathbf{Y}|\mathbf{X}, f) dp(f)} \end{aligned}$$

When is this well-defined?

# A posterior on functions?

How to find  $p(f|\mathbf{Y}, \mathbf{X})$ ?

*Bayes' rule:*

$$\begin{aligned} p(f|\mathbf{X}, \mathbf{Y}) &= \frac{p(\mathbf{Y}|\mathbf{X}, f) \cdot p(f)}{p(\mathbf{Y}|\mathbf{X})} \\ &= \frac{p(\mathbf{Y}|\mathbf{X}, f) \cdot p(f)}{\int p(\mathbf{Y}|\mathbf{X}, f) dp(f)} \end{aligned}$$

When is this well-defined?

# A posterior on functions?

Functions vs. parameters:

$$\mathcal{H} \cong \mathbb{R}^p$$

Represent functions in  $\mathcal{H}$  by their coordinates w.r.t. a basis:

$$f \in \mathcal{H} \leftrightarrow \theta \in \mathbb{R}^p$$

Assume (for the moment):  $p < \infty$

# A posterior on functions?

Functions vs. parameters:

$$\mathcal{H} \cong \mathbb{R}^p$$

Represent functions in  $\mathcal{H}$  by their coordinates w.r.t. a basis:

$$f \in \mathcal{H} \leftrightarrow \theta \in \mathbb{R}^p$$

Assume (for the moment):  $p < \infty$



# A posterior on functions?

*Mercer's theorem:*

$$K(\mathbf{x}_i, \mathbf{x}_j) = \sum_k \nu_k \psi_k(\mathbf{x}_i) \psi_k(\mathbf{x}_j)$$

where  $\nu_k \psi_k(\cdot) = \int K(\cdot, \mathbf{y}) \psi_k(\mathbf{y}) d\mathbf{y}$  for all  $k$ . The functions  $\{\sqrt{\nu_k} \psi_k(\cdot)\}$  form an *orthonormal basis* for  $\mathcal{H}_K$ .

Let  $\phi(\cdot) = [\sqrt{\nu_1} \psi_1(\cdot), \dots, \sqrt{\nu_p} \psi_p(\cdot)]$ . Then:

$$\mathcal{H}_K = \{\phi(\cdot)\theta \mid \theta \in \mathbb{R}^p\}$$

Problem: there's no such thing as

$$\theta \sim \mathcal{N}(0, I)$$

when  $\theta \in \mathbb{R}^\infty$ !

# Posterior for linear RLS

Linear function:

$$f(\mathbf{x}) = \langle \mathbf{x}, \theta \rangle$$

Noise model:

$$\mathbf{Y} | \mathbf{X}, \theta \sim \mathcal{N}(\mathbf{X}\theta, \sigma_\varepsilon^2 \mathbf{I})$$

Add a *prior*:

$$\theta \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

# Posterior for linear RLS

Model:

$$\mathbf{Y}|\mathbf{X}, \theta \sim \mathcal{N}(\mathbf{X}\theta, \sigma_\varepsilon^2 I), \quad \theta \sim \mathcal{N}(\mathbf{0}, I)$$

Joint over  $\mathbf{Y}$  and  $\theta$ :

$$\begin{bmatrix} \mathbf{Y} \\ \theta \end{bmatrix} \sim \mathcal{N}\left(\begin{bmatrix} \mathbf{0} \\ \mathbf{0} \end{bmatrix}, \begin{bmatrix} \mathbf{X}\mathbf{X}^T + \sigma_\varepsilon^2 I & \mathbf{X} \\ \mathbf{X}^T & I \end{bmatrix}\right)$$

Condition on  $\mathbf{Y}$ .

# Posterior for linear RLS

Posterior:

$$\theta|\mathbf{X}, \mathbf{Y} \sim \mathcal{N}(\mu_{\theta|\mathbf{X}, \mathbf{Y}}, \Sigma_{\theta|\mathbf{X}, \mathbf{Y}})$$

where

$$\mu_{\theta|\mathbf{X}, \mathbf{Y}} = \mathbf{X}^T(\mathbf{X}\mathbf{X}^T + \sigma_\varepsilon^2 I)^{-1}\mathbf{Y}$$

$$\Sigma_{\theta|\mathbf{X}, \mathbf{Y}} = I - \mathbf{X}^T(\mathbf{X}\mathbf{X}^T + \sigma_\varepsilon^2 I)^{-1}\mathbf{X}$$

This is Gaussian, so

$$\hat{\theta}_{MAP}(\mathbf{Y}|\mathbf{X}) = \hat{\theta}_{BLS}(\mathbf{Y}|\mathbf{X}) = \mathbf{X}^T(\mathbf{X}\mathbf{X}^T + \sigma_\varepsilon^2 I)^{-1}\mathbf{Y}$$

# Posterior for linear RLS

Posterior:

$$\theta|\mathbf{X}, \mathbf{Y} \sim \mathcal{N}(\mu_{\theta|\mathbf{X}, \mathbf{Y}}, \Sigma_{\theta|\mathbf{X}, \mathbf{Y}})$$

where

$$\mu_{\theta|\mathbf{X}, \mathbf{Y}} = \mathbf{X}^T (\mathbf{X}\mathbf{X}^T + \sigma_\varepsilon^2 I)^{-1} \mathbf{Y}$$

$$\Sigma_{\theta|\mathbf{X}, \mathbf{Y}} = I - \mathbf{X}^T (\mathbf{X}\mathbf{X}^T + \sigma_\varepsilon^2 I)^{-1} \mathbf{X}$$

This is Gaussian, so

$$\hat{\theta}_{MAP}(\mathbf{Y}|\mathbf{X}) = \hat{\theta}_{BLS}(\mathbf{Y}|\mathbf{X}) = \mathbf{X}^T (\mathbf{X}\mathbf{X}^T + \sigma_\varepsilon^2 I)^{-1} \mathbf{Y}$$

# Linear RLS as a MAP estimator

Model:

$$\mathbf{Y}|\mathbf{X}, \theta \sim \mathcal{N}(\mathbf{X}\theta, \sigma_\varepsilon^2 I), \quad \theta \sim \mathcal{N}(\mathbf{0}, I)$$

$$\hat{\theta}_{MAP}(\mathbf{Y}|\mathbf{X}) = \mathbf{X}^T(\mathbf{X}\mathbf{X}^T + \sigma_\varepsilon^2 I)^{-1}\mathbf{Y}$$

Recall the linear RLS solution:

$$\begin{aligned}\hat{\theta}_{RLS}(\mathbf{Y}|\mathbf{X}) &= \arg \min_{\theta} \frac{1}{2} \sum_{i=1}^N (y_i - \langle \mathbf{x}_i, \theta \rangle)^2 + \frac{\lambda}{2} \|\theta\|^2 \\ &= \mathbf{X}^T(\mathbf{X}\mathbf{X}^T + \frac{\lambda}{2} I)^{-1}\mathbf{Y}\end{aligned}$$

*So what's  $\lambda$ ?*

# Linear RLS as a MAP estimator

Model:

$$\mathbf{Y}|\mathbf{X}, \theta \sim \mathcal{N}(\mathbf{X}\theta, \sigma_\varepsilon^2 I), \quad \theta \sim \mathcal{N}(\mathbf{0}, I)$$

$$\hat{\theta}_{MAP}(\mathbf{Y}|\mathbf{X}) = \mathbf{X}^T(\mathbf{X}\mathbf{X}^T + \sigma_\varepsilon^2 I)^{-1}\mathbf{Y}$$

Recall the linear RLS solution:

$$\begin{aligned}\hat{\theta}_{RLS}(\mathbf{Y}|\mathbf{X}) &= \arg \min_{\theta} \frac{1}{2} \sum_{i=1}^N (y_i - \langle \mathbf{x}_i, \theta \rangle)^2 + \frac{\lambda}{2} \|\theta\|^2 \\ &= \mathbf{X}^T(\mathbf{X}\mathbf{X}^T + \frac{\lambda}{2} I)^{-1}\mathbf{Y}\end{aligned}$$

*So what's  $\lambda$ ?*



# Posterior for kernel RLS

Model for *linear* RLS:

$$\mathbf{Y}|\mathbf{X}, \theta \sim \mathcal{N}(\mathbf{X}\theta, \sigma_\varepsilon^2 \mathbf{I}), \quad \theta \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

Model for kernel RLS?

$$\mathbf{Y}|\mathbf{X}, \theta \sim \mathcal{N}(\phi(\mathbf{X})\theta, \sigma_\varepsilon^2 \mathbf{I}), \quad \theta \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

Then:

$$\hat{\theta}_{MAP}(\mathbf{Y}|\mathbf{X}) = \phi(\mathbf{X})^T (\phi(\mathbf{X})\phi(\mathbf{X})^T + \sigma_\varepsilon^2 \mathbf{I})^{-1} \mathbf{Y}$$

# Posterior for kernel RLS

Model for *linear* RLS:

$$\mathbf{Y}|\mathbf{X}, \theta \sim \mathcal{N}(\mathbf{X}\theta, \sigma_\varepsilon^2 \mathbf{I}), \quad \theta \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

Model for kernel RLS?

$$\mathbf{Y}|\mathbf{X}, \theta \sim \mathcal{N}(\phi(\mathbf{X})\theta, \sigma_\varepsilon^2 \mathbf{I}), \quad \theta \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

Then:

$$\hat{\theta}_{MAP}(\mathbf{Y}|\mathbf{X}) = \phi(\mathbf{X})^T (\phi(\mathbf{X})\phi(\mathbf{X})^T + \sigma_\varepsilon^2 \mathbf{I})^{-1} \mathbf{Y}$$

# Posterior for kernel RLS

Model for *linear* RLS:

$$\mathbf{Y}|\mathbf{X}, \theta \sim \mathcal{N}(\mathbf{X}\theta, \sigma_\varepsilon^2 \mathbf{I}), \quad \theta \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

Model for kernel RLS?

$$\mathbf{Y}|\mathbf{X}, \theta \sim \mathcal{N}(\phi(\mathbf{X})\theta, \sigma_\varepsilon^2 \mathbf{I}), \quad \theta \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

Then:

$$\hat{\theta}_{MAP}(\mathbf{Y}|\mathbf{X}) = \phi(\mathbf{X})^T (\mathbf{K} + \sigma_\varepsilon^2 \mathbf{I})^{-1} \mathbf{Y}$$

- **Empirical risk minimization is ML.**

$$p(\mathbf{Y}|f, \mathbf{X}) \propto e^{-\frac{1}{2} \sum_{i=1}^N (y_i - f(x_i))^2}$$

- Linear RLS is MAP.

$$p(\mathbf{Y}, f|\mathbf{X}) \propto e^{-\frac{1}{2} \sum_{i=1}^N (y_i - \langle x_i, \theta \rangle)^2} \cdot e^{-\frac{\lambda}{2} \theta^T \theta}$$

- Kernel RLS is also MAP.

$$p(\mathbf{Y}, f|\mathbf{X}) \propto e^{-\frac{1}{2} \sum_{i=1}^N (y_i - f(x_i))^2} \cdot e^{-\frac{\lambda}{2} \|f\|_{\mathcal{H}}^2}$$

- **Empirical risk minimization is ML.**

$$p(\mathbf{Y}|f, \mathbf{X}) \propto e^{-\frac{1}{2} \sum_{i=1}^N (y_i - f(x_i))^2}$$

- **Linear RLS is MAP.**

$$p(\mathbf{Y}, f|\mathbf{X}) \propto e^{-\frac{1}{2} \sum_{i=1}^N (y_i - \langle \mathbf{x}_i, \theta \rangle)^2} \cdot e^{-\frac{\lambda}{2} \theta^T \theta}$$

- **Kernel RLS is also MAP.**

$$p(\mathbf{Y}, f|\mathbf{X}) \propto e^{-\frac{1}{2} \sum_{i=1}^N (y_i - f(x_i))^2} \cdot e^{-\frac{\lambda}{2} \|f\|_{\mathcal{H}}^2}$$

- **Empirical risk minimization is ML.**

$$p(\mathbf{Y}|f, \mathbf{X}) \propto e^{-\frac{1}{2} \sum_{i=1}^N (y_i - f(x_i))^2}$$

- **Linear RLS is MAP.**

$$p(\mathbf{Y}, f|\mathbf{X}) \propto e^{-\frac{1}{2} \sum_{i=1}^N (y_i - \langle \mathbf{x}_i, \theta \rangle)^2} \cdot e^{-\frac{\lambda}{2} \theta^T \theta}$$

- **Kernel RLS is also MAP.**

$$p(\mathbf{Y}, f|\mathbf{X}) \propto e^{-\frac{1}{2} \sum_{i=1}^N (y_i - f(x_i))^2} \cdot e^{-\frac{\lambda}{2} \|f\|_{\mathcal{H}}^2}$$

# Transductive setting

*Idea:* Forget about estimating  $\theta$  (i.e.  $f$ ).

Instead: Estimate *predicted outputs*

$$\mathbf{Y}^* = [y_1^*, \dots, y_M^*]^T$$

at test inputs

$$\mathbf{X}^* = [x_1^*, \dots, x_M^*]^T$$

Need the joint distribution over  $\mathbf{Y}^*$  and  $\mathbf{Y}$ .

# Transductive setting

Say  $\mathbf{Y}^*$  and  $\mathbf{Y}$  are *jointly Gaussian*:

$$\begin{bmatrix} \mathbf{Y} \\ \mathbf{Y}^* \end{bmatrix} = \mathcal{N} \left( \begin{bmatrix} \mathbf{0} \\ \mathbf{0} \end{bmatrix}, \begin{bmatrix} \Lambda_{\mathbf{Y}} & \Lambda_{\mathbf{Y}\mathbf{Y}^*} \\ \Lambda_{\mathbf{Y}^*\mathbf{Y}} & \Lambda_{\mathbf{Y}^*} \end{bmatrix} \right)$$

Want: kernel RLS.

General form for the posterior:

$$\mathbf{Y}^* | \mathbf{X}, \mathbf{Y} \sim \mathcal{N}(\mu_{\mathbf{Y}^* | \mathbf{X}, \mathbf{Y}}, \Sigma_{\mathbf{Y}^* | \mathbf{X}, \mathbf{Y}})$$

where

$$\begin{aligned} \mu_{\mathbf{Y}^* | \mathbf{X}, \mathbf{Y}} &= \Lambda_{\mathbf{Y}\mathbf{Y}^*}^T \Lambda_{\mathbf{Y}}^{-1} \mathbf{Y} \\ \Sigma_{\mathbf{Y}^* | \mathbf{X}, \mathbf{Y}} &= \Lambda_{\mathbf{Y}^*} - \Lambda_{\mathbf{Y}\mathbf{Y}^*}^T \Lambda_{\mathbf{Y}}^{-1} \Lambda_{\mathbf{Y}\mathbf{Y}^*} \end{aligned}$$



# Transductive setting

Say  $\mathbf{Y}^*$  and  $\mathbf{Y}$  are *jointly Gaussian*:

$$\begin{bmatrix} \mathbf{Y} \\ \mathbf{Y}^* \end{bmatrix} = \mathcal{N} \left( \begin{bmatrix} \mathbf{0} \\ \mathbf{0} \end{bmatrix}, \begin{bmatrix} \Lambda_{\mathbf{Y}} & \Lambda_{\mathbf{Y}\mathbf{Y}^*} \\ \Lambda_{\mathbf{Y}^*\mathbf{Y}} & \Lambda_{\mathbf{Y}^*} \end{bmatrix} \right)$$

Want: kernel RLS.

General form for the posterior:

$$\mathbf{Y}^* | \mathbf{X}, \mathbf{Y} \sim \mathcal{N}(\mu_{\mathbf{Y}^* | \mathbf{X}, \mathbf{Y}}, \Sigma_{\mathbf{Y}^* | \mathbf{X}, \mathbf{Y}})$$

where

$$\mu_{\mathbf{Y}^* | \mathbf{X}, \mathbf{Y}} = \Lambda_{\mathbf{Y}\mathbf{Y}^*}^T \Lambda_{\mathbf{Y}}^{-1} \mathbf{Y}$$

$$\Sigma_{\mathbf{Y}^* | \mathbf{X}, \mathbf{Y}} = \Lambda_{\mathbf{Y}^*} - \Lambda_{\mathbf{Y}\mathbf{Y}^*}^T \Lambda_{\mathbf{Y}}^{-1} \Lambda_{\mathbf{Y}\mathbf{Y}^*}$$

$$\text{Set } \Lambda_{\mathbf{Y}} = K(\mathbf{X}, \mathbf{X}) + \sigma^2 I, \Lambda_{\mathbf{Y}Y^*} = K(\mathbf{X}, X^*), \Lambda_{Y^*} = K(X^*, X^*).$$

Posterior:

$$Y^* | \mathbf{X}, \mathbf{Y} \sim \mathcal{N}(\mu_{Y^* | \mathbf{X}, \mathbf{Y}}, \Sigma_{Y^* | \mathbf{X}, \mathbf{Y}})$$

where

$$\mu_{Y^* | \mathbf{X}, \mathbf{Y}} = K(X^*, \mathbf{X})(K(\mathbf{X}, \mathbf{X}) + \sigma^2 I)^{-1} \mathbf{Y}$$

$$\Sigma_{Y^* | \mathbf{X}, \mathbf{Y}} = K(X^*, X^*) - K(X^*, \mathbf{X})(K(\mathbf{X}, \mathbf{X}) + \sigma^2 I)^{-1} K(\mathbf{X}, X^*)$$

$$\text{So: } \hat{Y}_{MAP}^* = \hat{f}_{RLS}(X^*).$$

# Transductive setting

Model:

$$\begin{bmatrix} \mathbf{Y} \\ \mathbf{Y}^* \end{bmatrix} = \mathcal{N} \left( \begin{bmatrix} \mathbf{0} \\ \mathbf{0} \end{bmatrix}, \begin{bmatrix} \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_\varepsilon^2 \mathbf{I} & \mathbf{K}(\mathbf{X}, \mathbf{X}^*) \\ \mathbf{K}(\mathbf{X}^*, \mathbf{X}) & \mathbf{K}(\mathbf{X}^*, \mathbf{X}^*) \end{bmatrix} \right)$$

MAP estimate (posterior mean) = RLS function *at every point*  $x^*$ , regardless of  $\dim \mathcal{H}_K$ .

Are the prior and posterior (*on points!*) consistent with a distribution on  $\mathcal{H}_K$ ?

Strictly speaking,  $\theta$  and  $f$  don't come into play here at all:

Have:  $p(Y^*|\mathbf{X}, \mathbf{Y})$

Do not have:  $p(\theta|\mathbf{X}, \mathbf{Y})$  or  $p(f|\mathbf{X}, \mathbf{Y})$

But, *if  $\mathcal{H}_K$  is finite dimensional*, the joint over  $Y$  and  $Y^*$  is consistent with:

- $\mathbf{Y} = f(\mathbf{X}) + \varepsilon$ ,
- $Y^* = f(\mathbf{X})$ , and
- $f \in \mathcal{H}_K$  is a random trajectory from a **Gaussian process** over the domain, with mean  $\mu$  and covariance  $K$ .
- (Ergo, people call this “Gaussian process regression.”)  
(Also “Kriging,” because of a guy.)

Strictly speaking,  $\theta$  and  $f$  don't come into play here at all:

Have:  $p(Y^*|\mathbf{X}, \mathbf{Y})$

Do not have:  $p(\theta|\mathbf{X}, \mathbf{Y})$  or  $p(f|\mathbf{X}, \mathbf{Y})$

But, *if  $\mathcal{H}_K$  is finite dimensional*, the joint over  $\mathbf{Y}$  and  $Y^*$  is consistent with:

- $\mathbf{Y} = f(\mathbf{X}) + \varepsilon$ ,
- $Y^* = f(\mathbf{X})$ , and
- $f \in \mathcal{H}_K$  is a random trajectory from a **Gaussian process** over the domain, with mean  $\mu$  and covariance  $K$ .
- (Ergo, people call this “Gaussian process regression.”)  
(Also “Kriging,” because of a guy.)

- **Empirical risk minimization** is the maximum likelihood estimator when:

$$y = \mathbf{x}^T \theta + \varepsilon$$

- **Linear RLS** is the MAP estimator when:

$$y = \mathbf{x}^T \theta + \varepsilon, \quad \theta \sim \mathcal{N}(\mathbf{0}, I)$$

- **Kernel RLS** is the MAP estimator when:

$$y = \phi(\mathbf{x})^T \theta + \varepsilon, \quad \theta \sim \mathcal{N}(\mathbf{0}, I)$$

*in finite dimensional  $\mathcal{H}_K$ .*

- **Kernel RLS** is the MAP estimator *at points* when:

$$\begin{bmatrix} \mathbf{Y} \\ \mathbf{Y}^* \end{bmatrix} = \mathcal{N} \left( \begin{bmatrix} \mu_{\mathbf{Y}} \\ \mu_{\mathbf{Y}^*} \end{bmatrix}, \begin{bmatrix} \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_{\varepsilon}^2 I & \mathbf{K}(\mathbf{X}, \mathbf{X}^*) \\ \mathbf{K}(\mathbf{X}^*, \mathbf{X}) & \mathbf{K}(\mathbf{X}^*, \mathbf{X}^*) \end{bmatrix} \right)$$

*in possibly infinite dimensional  $\mathcal{H}_K$ .*

# Is this useful in practice?

- Want confidence intervals + believe the posteriors are meaningful = yes
- Maybe other reasons?

