

Functional Analysis Review

Lorenzo Rosasco
–slides courtesy of Andre Wibisono

9.520: Statistical Learning Theory and Applications

February 13, 2012

- 1 Vector Spaces
- 2 Hilbert Spaces
- 3 Matrices
- 4 Linear Operators

Vector Space

- A **vector space** is a set V with binary operations

$$+ : V \times V \rightarrow V \quad \text{and} \quad \cdot : \mathbb{R} \times V \rightarrow V$$

such that for all $a, b \in \mathbb{R}$ and $v, w, x \in V$:

- 1 $v + w = w + v$
- 2 $(v + w) + x = v + (w + x)$
- 3 There exists $0 \in V$ such that $v + 0 = v$ for all $v \in V$
- 4 For every $v \in V$ there exists $-v \in V$ such that $v + (-v) = 0$
- 5 $a(bv) = (ab)v$
- 6 $1v = v$
- 7 $(a + b)v = av + bv$
- 8 $a(v + w) = av + aw$

Vector Space

- A **vector space** is a set V with binary operations

$$+ : V \times V \rightarrow V \quad \text{and} \quad \cdot : \mathbb{R} \times V \rightarrow V$$

such that for all $a, b \in \mathbb{R}$ and $v, w, x \in V$:

- ① $v + w = w + v$
 - ② $(v + w) + x = v + (w + x)$
 - ③ There exists $0 \in V$ such that $v + 0 = v$ for all $v \in V$
 - ④ For every $v \in V$ there exists $-v \in V$ such that $v + (-v) = 0$
 - ⑤ $a(bv) = (ab)v$
 - ⑥ $1v = v$
 - ⑦ $(a + b)v = av + bv$
 - ⑧ $a(v + w) = av + aw$
- Example: \mathbb{R}^n , space of polynomials, space of functions.

Inner Product

- An **inner product** is a function $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$ such that for all $\alpha, \beta \in \mathbb{R}$ and $v, w, x \in V$:

Inner Product

- An **inner product** is a function $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$ such that for all $a, b \in \mathbb{R}$ and $v, w, x \in V$:
 - ① $\langle v, w \rangle = \langle w, v \rangle$
 - ② $\langle av + bw, x \rangle = a\langle v, x \rangle + b\langle w, x \rangle$
 - ③ $\langle v, v \rangle \geq 0$ and $\langle v, v \rangle = 0$ if and only if $v = 0$.

Inner Product

- An **inner product** is a function $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$ such that for all $a, b \in \mathbb{R}$ and $v, w, x \in V$:
 - 1 $\langle v, w \rangle = \langle w, v \rangle$
 - 2 $\langle av + bw, x \rangle = a\langle v, x \rangle + b\langle w, x \rangle$
 - 3 $\langle v, v \rangle \geq 0$ and $\langle v, v \rangle = 0$ if and only if $v = 0$.
- $v, w \in V$ are orthogonal if $\langle v, w \rangle = 0$.

Inner Product

- An **inner product** is a function $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$ such that for all $a, b \in \mathbb{R}$ and $v, w, x \in V$:
 - 1 $\langle v, w \rangle = \langle w, v \rangle$
 - 2 $\langle av + bw, x \rangle = a\langle v, x \rangle + b\langle w, x \rangle$
 - 3 $\langle v, v \rangle \geq 0$ and $\langle v, v \rangle = 0$ if and only if $v = 0$.
- $v, w \in V$ are orthogonal if $\langle v, w \rangle = 0$.
- Given $W \subseteq V$, we have $V = W \oplus W^\perp$, where $W^\perp = \{ v \in V \mid \langle v, w \rangle = 0 \text{ for all } w \in W \}$.

Inner Product

- An **inner product** is a function $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$ such that for all $a, b \in \mathbb{R}$ and $v, w, x \in V$:
 - 1 $\langle v, w \rangle = \langle w, v \rangle$
 - 2 $\langle av + bw, x \rangle = a\langle v, x \rangle + b\langle w, x \rangle$
 - 3 $\langle v, v \rangle \geq 0$ and $\langle v, v \rangle = 0$ if and only if $v = 0$.
- $v, w \in V$ are orthogonal if $\langle v, w \rangle = 0$.
- Given $W \subseteq V$, we have $V = W \oplus W^\perp$, where $W^\perp = \{ v \in V \mid \langle v, w \rangle = 0 \text{ for all } w \in W \}$.
- Cauchy-Schwarz inequality: $\langle v, w \rangle \leq \langle v, v \rangle^{1/2} \langle w, w \rangle^{1/2}$.

Norm

- Can define norm from inner product: $\|v\| = \langle v, v \rangle^{1/2}$.

Norm

- A **norm** is a function $\| \cdot \|: V \rightarrow \mathbb{R}$ such that for all $\alpha \in \mathbb{R}$ and $v, w \in V$:
 - 1 $\|v\| \geq 0$, and $\|v\| = 0$ if and only if $v = 0$
 - 2 $\|\alpha v\| = |\alpha| \|v\|$
 - 3 $\|v + w\| \leq \|v\| + \|w\|$
- Can define norm from inner product: $\|v\| = \langle v, v \rangle^{1/2}$.

Metric

- Can define metric from norm: $d(v, w) = \|v - w\|$.

Metric

- A **metric** is a function $d: V \times V \rightarrow \mathbb{R}$ such that for all $v, w, x \in V$:
 - ① $d(v, w) \geq 0$, and $d(v, w) = 0$ if and only if $v = w$
 - ② $d(v, w) = d(w, v)$
 - ③ $d(v, w) \leq d(v, x) + d(x, w)$
- Can define metric from norm: $d(v, w) = \|v - w\|$.

Basis

- $B = \{v_1, \dots, v_n\}$ is a **basis** of V if every $v \in V$ can be uniquely decomposed as

$$v = \alpha_1 v_1 + \dots + \alpha_n v_n$$

for some $\alpha_1, \dots, \alpha_n \in \mathbb{R}$.

Basis

- $B = \{v_1, \dots, v_n\}$ is a **basis** of V if every $v \in V$ can be uniquely decomposed as

$$v = a_1 v_1 + \dots + a_n v_n$$

for some $a_1, \dots, a_n \in \mathbb{R}$.

- An orthonormal basis is a basis that is orthogonal ($\langle v_i, v_j \rangle = 0$ for $i \neq j$) and normalized ($\|v_i\| = 1$).

- 1 Vector Spaces
- 2 Hilbert Spaces**
- 3 Matrices
- 4 Linear Operators

Hilbert Space, overview

- Goal: to understand Hilbert spaces (complete inner product spaces) and to make sense of the expression

$$f = \sum_{i=1}^{\infty} \langle f, \phi_i \rangle \phi_i, \quad f \in \mathcal{H}$$

- Need to talk about:
 - 1 Cauchy sequence
 - 2 Completeness
 - 3 Density
 - 4 Separability

Cauchy Sequence

- Recall: $\lim_{n \rightarrow \infty} x_n = x$ if for every $\epsilon > 0$ there exists $N \in \mathbb{N}$ such that $\|x - x_n\| < \epsilon$ whenever $n \geq N$.

Cauchy Sequence

- Recall: $\lim_{n \rightarrow \infty} x_n = x$ if for every $\epsilon > 0$ there exists $N \in \mathbb{N}$ such that $\|x - x_n\| < \epsilon$ whenever $n \geq N$.
- $(x_n)_{n \in \mathbb{N}}$ is a **Cauchy sequence** if for every $\epsilon > 0$ there exists $N \in \mathbb{N}$ such that $\|x_m - x_n\| < \epsilon$ whenever $m, n \geq N$.

Cauchy Sequence

- Recall: $\lim_{n \rightarrow \infty} x_n = x$ if for every $\epsilon > 0$ there exists $N \in \mathbb{N}$ such that $\|x - x_n\| < \epsilon$ whenever $n \geq N$.
- $(x_n)_{n \in \mathbb{N}}$ is a **Cauchy sequence** if for every $\epsilon > 0$ there exists $N \in \mathbb{N}$ such that $\|x_m - x_n\| < \epsilon$ whenever $m, n \geq N$.
- Every convergent sequence is a Cauchy sequence (why?)

Completeness

- A normed vector space V is **complete** if every Cauchy sequence converges.

Completeness

- A normed vector space V is **complete** if every Cauchy sequence converges.
- Examples:
 - 1 \mathbb{Q} is not complete.
 - 2 \mathbb{R} is complete (axiom).
 - 3 \mathbb{R}^n is complete.
 - 4 Every finite dimensional normed vector space (over \mathbb{R}) is complete.

Hilbert Space

- A **Hilbert space** is a complete inner product space.

Hilbert Space

- A **Hilbert space** is a complete inner product space.
- Examples:
 - ① \mathbb{R}^n
 - ② Every finite dimensional inner product space.
 - ③ $\ell_2 = \{(\mathbf{a}_n)_{n=1}^{\infty} \mid \mathbf{a}_n \in \mathbb{R}, \sum_{n=1}^{\infty} \mathbf{a}_n^2 < \infty\}$
 - ④ $L_2([0, 1]) = \{f: [0, 1] \rightarrow \mathbb{R} \mid \int_0^1 f(x)^2 dx < \infty\}$

Density

- Y is **dense** in X if $\overline{Y} = X$.

Density

- Y is **dense** in X if $\overline{Y} = X$.
- Examples:
 - 1 \mathbb{Q} is dense in \mathbb{R} .
 - 2 \mathbb{Q}^n is dense in \mathbb{R}^n .
 - 3 Weierstrass approximation theorem: polynomials are dense in continuous functions (with the supremum norm, on compact domains).

Separability

- X is **separable** if it has a countable dense subset.

Separability

- X is **separable** if it has a countable dense subset.
- Examples:
 - 1 \mathbb{R} is separable.
 - 2 \mathbb{R}^n is separable.
 - 3 ℓ_2 , $L_2([0, 1])$ are separable.

Orthonormal Basis

- A Hilbert space has a countable orthonormal basis if and only if it is separable.
- Can write:

$$f = \sum_{i=1}^{\infty} \langle f, \phi_i \rangle \phi_i \quad \text{for all } f \in \mathcal{H}.$$

Orthonormal Basis

- A Hilbert space has a countable orthonormal basis if and only if it is separable.
- Can write:

$$f = \sum_{i=1}^{\infty} \langle f, \phi_i \rangle \phi_i \quad \text{for all } f \in \mathcal{H}.$$

- Examples:
 - 1 Basis of ℓ_2 is $(1, 0, \dots), (0, 1, 0, \dots), (0, 0, 1, 0, \dots), \dots$
 - 2 Basis of $L_2([0, 1])$ is $1, 2 \sin 2\pi n x, 2 \cos 2\pi n x$ for $n \in \mathbb{N}$

Maps

Next we are going to review basic properties of maps on a Hilbert space.

- functionals: $\Psi : \mathcal{H} \rightarrow \mathbb{R}$
- linear operators $A : \mathcal{H} \rightarrow \mathcal{H}$, such that
 $A(af + bg) = aAf + bAg$, with $a, b \in \mathbb{R}$ and $f, g \in \mathcal{H}$.

Representation of Continuous Functionals

Let \mathcal{H} be a Hilbert space and $g \in \mathcal{H}$, then

$$\Psi_g(f) = \langle f, g \rangle, \quad f \in \mathcal{H}$$

is a continuous linear functional.

Riesz representation theorem

The theorem states that every continuous linear functional Ψ can be written uniquely in the form,

$$\Psi(f) = \langle f, g \rangle$$

for some appropriate element $g \in \mathcal{H}$.

- 1 Vector Spaces
- 2 Hilbert Spaces
- 3 Matrices**
- 4 Linear Operators

Matrix

- Every linear operator $L: \mathbb{R}^m \rightarrow \mathbb{R}^n$ can be represented by an $m \times n$ matrix A .

Matrix

- Every linear operator $L: \mathbb{R}^m \rightarrow \mathbb{R}^n$ can be represented by an $m \times n$ matrix A .
- If $A \in \mathbb{R}^{m \times n}$, the transpose of A is $A^T \in \mathbb{R}^{n \times m}$ satisfying

$$\langle Ax, y \rangle_{\mathbb{R}^m} = (Ax)^T y = x^T A^T y = \langle x, A^T y \rangle_{\mathbb{R}^n}$$

for every $x \in \mathbb{R}^n$ and $y \in \mathbb{R}^m$.

Matrix

- Every linear operator $L: \mathbb{R}^m \rightarrow \mathbb{R}^n$ can be represented by an $m \times n$ matrix A .
- If $A \in \mathbb{R}^{m \times n}$, the transpose of A is $A^T \in \mathbb{R}^{n \times m}$ satisfying

$$\langle Ax, y \rangle_{\mathbb{R}^m} = (Ax)^T y = x^T A^T y = \langle x, A^T y \rangle_{\mathbb{R}^n}$$

for every $x \in \mathbb{R}^n$ and $y \in \mathbb{R}^m$.

- A is symmetric if $A^T = A$.

Eigenvalues and Eigenvectors

- Let $A \in \mathbb{R}^{n \times n}$. A nonzero vector $v \in \mathbb{R}^n$ is an eigenvector of A with corresponding eigenvalue $\lambda \in \mathbb{R}$ if $Av = \lambda v$.

Eigenvalues and Eigenvectors

- Let $A \in \mathbb{R}^{n \times n}$. A nonzero vector $v \in \mathbb{R}^n$ is an eigenvector of A with corresponding eigenvalue $\lambda \in \mathbb{R}$ if $Av = \lambda v$.
- Symmetric matrices have real eigenvalues.

Eigenvalues and Eigenvectors

- Let $A \in \mathbb{R}^{n \times n}$. A nonzero vector $v \in \mathbb{R}^n$ is an eigenvector of A with corresponding eigenvalue $\lambda \in \mathbb{R}$ if $Av = \lambda v$.
- Symmetric matrices have real eigenvalues.
- **Spectral Theorem:** Let A be a symmetric $n \times n$ matrix. Then there is an orthonormal basis of \mathbb{R}^n consisting of the eigenvectors of A .

Eigenvalues and Eigenvectors

- Let $A \in \mathbb{R}^{n \times n}$. A nonzero vector $v \in \mathbb{R}^n$ is an eigenvector of A with corresponding eigenvalue $\lambda \in \mathbb{R}$ if $Av = \lambda v$.
- Symmetric matrices have real eigenvalues.
- **Spectral Theorem:** Let A be a symmetric $n \times n$ matrix. Then there is an orthonormal basis of \mathbb{R}^n consisting of the eigenvectors of A .
- Eigendecomposition: $A = V\Lambda V^T$, or equivalently,

$$A = \sum_{i=1}^n \lambda_i v_i v_i^T.$$

Singular Value Decomposition

- Every $A \in \mathbb{R}^{m \times n}$ can be written as

$$A = U\Sigma V^T,$$

where $U \in \mathbb{R}^{m \times m}$ is orthogonal, $\Sigma \in \mathbb{R}^{m \times n}$ is diagonal, and $V \in \mathbb{R}^{n \times n}$ is orthogonal.

Singular Value Decomposition

- Every $A \in \mathbb{R}^{m \times n}$ can be written as

$$A = U\Sigma V^T,$$

where $U \in \mathbb{R}^{m \times m}$ is orthogonal, $\Sigma \in \mathbb{R}^{m \times n}$ is diagonal, and $V \in \mathbb{R}^{n \times n}$ is orthogonal.

- Singular system:

$$\begin{aligned} Av_i &= \sigma_i u_i & AA^T u_i &= \sigma_i^2 u_i \\ A^T u_i &= \sigma_i v_i & A^T Av_i &= \sigma_i^2 v_i \end{aligned}$$

Matrix Norm

- The spectral norm of $\mathbf{A} \in \mathbb{R}^{m \times n}$ is

$$\|\mathbf{A}\|_{\text{spec}} = \sigma_{\max}(\mathbf{A}) = \sqrt{\lambda_{\max}(\mathbf{A}\mathbf{A}^{\top})} = \sqrt{\lambda_{\max}(\mathbf{A}^{\top}\mathbf{A})}.$$

Matrix Norm

- The spectral norm of $\mathbf{A} \in \mathbb{R}^{m \times n}$ is

$$\|\mathbf{A}\|_{\text{spec}} = \sigma_{\max}(\mathbf{A}) = \sqrt{\lambda_{\max}(\mathbf{A}\mathbf{A}^{\top})} = \sqrt{\lambda_{\max}(\mathbf{A}^{\top}\mathbf{A})}.$$

- The Frobenius norm of $\mathbf{A} \in \mathbb{R}^{m \times n}$ is

$$\|\mathbf{A}\|_{\text{F}} = \sqrt{\sum_{i=1}^m \sum_{j=1}^n a_{ij}^2} = \sqrt{\sum_{i=1}^{\min\{m,n\}} \sigma_i^2}.$$

Positive Definite Matrix

A real symmetric matrix $A \in \mathbb{R}^{m \times m}$ is positive definite if

$$x^t A x > 0, \quad \forall x \in \mathbb{R}^m.$$

A positive definite matrix has positive eigenvalues.

Note: for positive semi-definite matrices $>$ is replaced by \geq .

- 1 Vector Spaces
- 2 Hilbert Spaces
- 3 Matrices
- 4 Linear Operators**

Linear Operator

- An operator $L: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is linear if it preserves the linear structure.

Linear Operator

- An operator $L: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is linear if it preserves the linear structure.
- A linear operator $L: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is bounded if there exists $C > 0$ such that

$$\|Lf\|_{\mathcal{H}_2} \leq C\|f\|_{\mathcal{H}_1} \quad \text{for all } f \in \mathcal{H}_1.$$

Linear Operator

- An operator $L: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is linear if it preserves the linear structure.
- A linear operator $L: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is bounded if there exists $C > 0$ such that

$$\|Lf\|_{\mathcal{H}_2} \leq C\|f\|_{\mathcal{H}_1} \quad \text{for all } f \in \mathcal{H}_1.$$

- A linear operator is continuous if and only if it is bounded.

Adjoint and Compactness

- The adjoint of a bounded linear operator $L: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is a bounded linear operator $L^*: \mathcal{H}_2 \rightarrow \mathcal{H}_1$ satisfying

$$\langle Lf, g \rangle_{\mathcal{H}_2} = \langle f, L^*g \rangle_{\mathcal{H}_1} \quad \text{for all } f \in \mathcal{H}_1, g \in \mathcal{H}_2.$$

- L is self-adjoint if $L^* = L$. Self-adjoint operators have real eigenvalues.

Adjoint and Compactness

- The adjoint of a bounded linear operator $L: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is a bounded linear operator $L^*: \mathcal{H}_2 \rightarrow \mathcal{H}_1$ satisfying

$$\langle Lf, g \rangle_{\mathcal{H}_2} = \langle f, L^*g \rangle_{\mathcal{H}_1} \quad \text{for all } f \in \mathcal{H}_1, g \in \mathcal{H}_2.$$

- L is self-adjoint if $L^* = L$. Self-adjoint operators have real eigenvalues.
- A bounded linear operator $L: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is compact if the image of the unit ball in \mathcal{H}_1 has compact closure in \mathcal{H}_2 .

Spectral Theorem for Compact Self-Adjoint Operator

- Let $L: \mathcal{H} \rightarrow \mathcal{H}$ be a compact self-adjoint operator. Then there exists an orthonormal basis of \mathcal{H} consisting of the eigenfunctions of L ,

$$L\phi_i = \lambda_i\phi_i$$

and the only possible limit point of λ_i as $i \rightarrow \infty$ is 0.

Spectral Theorem for Compact Self-Adjoint Operator

- Let $L: \mathcal{H} \rightarrow \mathcal{H}$ be a compact self-adjoint operator. Then there exists an orthonormal basis of \mathcal{H} consisting of the eigenfunctions of L ,

$$L\phi_i = \lambda_i \phi_i$$

and the only possible limit point of λ_i as $i \rightarrow \infty$ is 0.

- Eigendecomposition:

$$L = \sum_{i=1}^{\infty} \lambda_i \langle \phi_i, \cdot \rangle \phi_i.$$